

G, G<sup>1</sup> and G<sup>2</sup> are taken separately and are each hydrogen and R<sup>6</sup> is hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl or phenyl optionally independently substituted with up to three hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl or (C<sub>1</sub>-C<sub>4</sub>)alkoxy, wherein said  
5 (C<sub>1</sub>-C<sub>4</sub>)alkyl in the definition of R<sup>6</sup> and said (C<sub>1</sub>-C<sub>4</sub>)alkoxy in the definition of R<sup>6</sup> are optionally and independently substituted with up to five fluoro; R<sup>7</sup> and R<sup>8</sup> are each independently hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl; or

G and G<sup>1</sup> are taken together and are (C<sub>1</sub>-C<sub>3</sub>)alkylene and R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup> and G<sup>2</sup> are hydrogen; or

10 G<sup>1</sup> and G<sup>2</sup> are taken together and are (C<sub>1</sub>-C<sub>3</sub>)alkylene and R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup> and G are hydrogen;

q is 0 or 1;

X is a covalent bond, -(C=NR<sup>10</sup>)-, oxycarbonyl, vinylenylcarbonyl, oxy(C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl, (C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl, (C<sub>3</sub>-C<sub>4</sub>)alkenylcarbonyl, thio(C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl, vinylenylsulfonyl, sulfinyl-(C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl, sulfonyl-(C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl or carbonyl(C<sub>0</sub>-C<sub>4</sub>)alkylenylcarbonyl; wherein said oxy(C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl, (C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl, (C<sub>3</sub>-C<sub>4</sub>)alkenylcarbonyl and thio(C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl in the definition of X are each optionally and independently substituted with up to two (C<sub>1</sub>-C<sub>4</sub>)alkyl, benzyl or Ar; said vinylenylsulfonyl and said  
20 vinylenylcarbonyl in the definition of X are optionally substituted independently on one or two vinylenyl carbons with (C<sub>1</sub>-C<sub>4</sub>)alkyl, benzyl or Ar; and said carbonyl(C<sub>0</sub>-C<sub>4</sub>)alkylenylcarbonyl in the definition of X is optionally substituted independently with up to three (C<sub>1</sub>-C<sub>4</sub>)alkyl, benzyl or Ar;

R<sup>10</sup> is hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl;

25 R<sup>9</sup> is (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, Ar<sup>1</sup>-(C<sub>0</sub>-C<sub>3</sub>)alkylenyl or (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted with up to five fluoro; provided that when q = 0 and X is a covalent bond, oxycarbonyl or (C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl, then R<sup>9</sup> is not (C<sub>1</sub>-C<sub>6</sub>)alkyl;

Ar and Ar<sup>1</sup> are independently a fully saturated, partially saturated or fully unsaturated five- to eight-membered ring optionally having up to four heteroatoms selected  
30 independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of two fused independently partially saturated, fully saturated or fully unsaturated five- to seven-membered rings, taken independently, optionally having up to four heteroatoms selected independently from nitrogen, sulfur and oxygen, or a tricyclic ring consisting of three fused independently partially saturated, fully saturated or fully  
35 unsaturated five to seven membered rings, taken independently, optionally having up

to four heteroatoms selected independently from nitrogen, sulfur and oxygen, said partially saturated, fully saturated ring or fully unsaturated monocyclic ring, bicyclic ring or tricyclic ring optionally having one or two oxo groups substituted on carbon or one or two oxo groups substituted on sulfur;

- 5 Ar and Ar<sup>1</sup> are optionally independently substituted on carbon or nitrogen, on one ring if the moiety is monocyclic, on one or both rings if the moiety is bicyclic, or on one, two or three rings if the moiety is tricyclic, with up to a total of four substituents independently selected from R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup>; wherein R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> are each taken separately and are each independently halo, formyl, (C<sub>1</sub>-
- 10 C<sub>6</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylenyloxycarbonyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, C(OH)R<sup>15</sup>R<sup>16</sup>, naphthyl, phenyl, imidazolyl, pyridyl, triazolyl, morpholinyl, (C<sub>0</sub>-C<sub>4</sub>)alkylsulfamoyl, N-(C<sub>0</sub>-C<sub>4</sub>)alkylcarbamoyl, N,N-di-(C<sub>1</sub>-C<sub>4</sub>)alkylcarbamoyl, N-phenylcarbamoyl, N-(C<sub>1</sub>-C<sub>4</sub>)alkyl-N-phenylcarbamoyl, N,N-diphenyl carbamoyl, (C<sub>1</sub>-C<sub>4</sub>)alkylcarbonylamido, (C<sub>3</sub>-C<sub>7</sub>)cycloalkylcarbonylamido, phenylcarbonylamido,
- 15 piperidinyl, pyrrolidinyl, piperazinyl, cyano, benzimidazolyl, amino, anilino, pyrimidyl, oxazolyl, isoxazolyl, tetrazolyl, thienyl, thiazolyl, benzothiazolyl, pyrrolyl, pyrazolyl, tetrahydroquinolyl, tetrahydroisoquinolyl, benzoxazolyl, pyridazinyl, pyridyloxy, pyridylsulfanyl, furanyl, 8-(C<sub>1</sub>-C<sub>4</sub>)alkyl-3,8-diaza[3.2.1]bicyclooctyl, 3,5-dioxo-1,2,4-triazinyl, phenoxy, thiophenoxy, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfanyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonyl, (C<sub>3</sub>-
- 20 C<sub>7</sub>)cycloalkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said naphthyl, phenyl, pyridyl, piperidinyl, benzimidazolyl, pyrimidyl, thienyl, benzothiazolyl, pyrrolyl, tetrahydroquinolyl, tetrahydroisoquinolyl, benzoxazolyl, pyridazinyl, pyridyloxy, pyridylsulfanyl, furanyl, thiophenoxy, anilino and phenoxy in the definition of R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> are
- 25 optionally substituted with up to three substituents independently selected from hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said imidazolyl, oxazolyl, isoxazolyl, thiazolyl and pyrazolyl in the definition of R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> are optionally substituted with up to two substituents
- 30 independently selected from hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said morpholinyl in the definition of R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> is optionally substituted with up to two substituents independently selected from (C<sub>1</sub>-C<sub>4</sub>)alkyl; said pyrrolidinyl in the definition of R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> is
- 35 optionally substituted with up to two substituents independently selected from

hydroxy, hydroxy-(C<sub>1</sub>-C<sub>3</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said piperazinyl in the definition of R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> is optionally substituted with up to three substituents independently selected from (C<sub>1</sub>-C<sub>4</sub>)alkoxy-  
5 (C<sub>1</sub>-C<sub>4</sub>)alkyl, hydroxy-(C<sub>1</sub>-C<sub>3</sub>)alkyl, phenyl, pyridyl, (C<sub>0</sub>-C<sub>4</sub>)alkylsulfamoyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said triazolyl in the definition of R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> is optionally substituted with hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy  
10 optionally substituted with up to five fluoro; said tetrazolyl in the definition of R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> is optionally substituted with hydroxy-(C<sub>2</sub>-C<sub>3</sub>)alkyl or (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro; and said phenyl and pyridyl which are optionally substituted on piperazine in the definition of R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> are optionally substituted with up to three hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-  
15 C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; or R<sup>11</sup> and R<sup>12</sup> are taken together on adjacent carbon atoms and are -CH<sub>2</sub>OC(CH<sub>3</sub>)<sub>2</sub>OCH<sub>2</sub>- or -O-(CH<sub>2</sub>)<sub>p</sub>-O-, and R<sup>13</sup> and R<sup>14</sup> are taken separately and are each independently hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl;  
20 p is 1, 2 or 3; R<sup>15</sup> and R<sup>16</sup> are taken separately and are each independently hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro; or R<sup>15</sup> and R<sup>16</sup> are taken separately and R<sup>15</sup> is hydrogen and R<sup>16</sup> is (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, hydroxy-(C<sub>1</sub>-C<sub>3</sub>)alkyl, phenyl, pyridyl, pyrimidyl, thienyl, furanyl, thiazolyl, oxazolyl, imidazolyl, benzothiazolyl or  
25 benzoxazolyl; or R<sup>15</sup> and R<sup>16</sup> are taken together and are (C<sub>3</sub>-C<sub>6</sub>)alkylene; G<sup>3</sup>, G<sup>4</sup> and G<sup>5</sup> are taken separately and are each hydrogen; r is 0; R<sup>18</sup> is hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl or phenyl optionally independently substituted with up to three hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl or (C<sub>1</sub>-C<sub>4</sub>)alkoxy,  
30 wherein said (C<sub>1</sub>-C<sub>4</sub>)alkyl in the definition of R<sup>6</sup> and said (C<sub>1</sub>-C<sub>4</sub>)alkoxy in the definition of R<sup>6</sup> are optionally and independently substituted with up to five fluoro; and R<sup>19</sup> and R<sup>20</sup> are each independently (C<sub>1</sub>-C<sub>4</sub>)alkyl; or G<sup>3</sup>, G<sup>4</sup> and G<sup>5</sup> are taken separately and are each hydrogen; r is 1; R<sup>18</sup> is hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl  
35 or phenyl optionally independently substituted with up to three hydroxy, halo,

hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl or (C<sub>1</sub>-C<sub>4</sub>)alkoxy, wherein said (C<sub>1</sub>-C<sub>4</sub>)alkyl in the definition of R<sup>6</sup> and said (C<sub>1</sub>-C<sub>4</sub>)alkoxy in the definition of R<sup>6</sup> are optionally and independently substituted with up to five fluoro; and R<sup>19</sup> and R<sup>20</sup> are each independently hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl; or

5 G<sup>3</sup> and G<sup>4</sup> are taken together and are (C<sub>1</sub>-C<sub>3</sub>)alkylene; r is 0 or 1; and R<sup>18</sup>, R<sup>19</sup>, R<sup>20</sup> and G<sup>5</sup> are hydrogen; or

G<sup>4</sup> and G<sup>5</sup> are taken together and are (C<sub>1</sub>-C<sub>3</sub>)alkylene; r is 0 or 1; and R<sup>18</sup>, R<sup>19</sup>, R<sup>20</sup> and G<sup>3</sup> are hydrogen;

10 R<sup>17</sup> is SO<sub>2</sub>NR<sup>21</sup>R<sup>22</sup>, CONR<sup>21</sup>R<sup>22</sup>, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl, Ar<sup>2</sup>-carbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfinyl, Ar<sup>2</sup>-sulfonyl, Ar<sup>2</sup>-sufinyl and (C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sup>21</sup> and R<sup>22</sup> are taken separately and are each independently selected from hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl and Ar<sup>2</sup>-(C<sub>0</sub>-C<sub>4</sub>)alkylenyl; or

R<sup>21</sup> and R<sup>22</sup> are taken together with the nitrogen atom to which they are attached to

15 form azetidiny, pyrrolidiny, piperidiny, piperaziny, morpholiny, azepiny, azabicyclo[3.2.2]nonany, azabicyclo[2.2.1]heptyl, 6,7-dihydro-5H-dibenzo[c,e]azepiny, 1,2,3,4-tetrahydro-isoquinoly or 5,6,7,8-tetrahydropyrido[4,3-d]pyrimidy; said azetidiny in the definition of R<sup>21</sup> and R<sup>22</sup> is optionally substituted independently with one substituent selected from hydroxy, amino, hydroxy-(C<sub>1</sub>-

20 C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said pyrrolidiny, piperidiny, azepiny in the definition of R<sup>21</sup> and R<sup>22</sup> are optionally substituted independently with up to two substituents independently selected from hydroxy, amino, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally

25 substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said morpholiny in the definition of R<sup>21</sup> and R<sup>22</sup> is optionally substituted with up to two substituents independently selected from hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said piperaziny in the

30 definition of R<sup>21</sup> and R<sup>22</sup> is optionally substituted independently with up to three substituents independently selected from phenyl, pyridyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl and (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro; said 1,2,3,4-tetrahydro-isoquinoly and said 5,6,7,8-tetrahydropyrido[4,3-d]pyrimidy in the definition of R<sup>21</sup> and R<sup>22</sup> are optionally substituted independently with up to three substituents

35 independently selected from hydroxy, amino, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-

C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; and said 6,7-dihydro-5H-dibenzo[c,e]azepinyl in the definition of R<sup>21</sup> and R<sup>22</sup> is optionally substituted with up to four substituents independently selected from hydroxy, amino, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said pyrimidyl, pyridyl and phenyl which are optionally substituted on said piperazine in the definition of R<sup>21</sup> and R<sup>22</sup> is optionally substituted with up to three substituents selected from hydroxy, amino, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro;

Ar<sup>2</sup> is independently defined as set forth for Ar and Ar<sup>1</sup> above;  
said Ar<sup>2</sup> is optionally independently substituted as set forth for Ar and Ar<sup>1</sup> above;  
R<sup>23</sup> is CONR<sup>25</sup>R<sup>26</sup> or SO<sub>2</sub>R<sup>25</sup>R<sup>26</sup>, wherein R<sup>25</sup> is hydrogen (C<sub>1</sub>-C<sub>4</sub>)alkyl or Ar<sup>3</sup>-(C<sub>0</sub>-C<sub>4</sub>)alkylenyl and R<sup>26</sup> is Ar<sup>3</sup>-(C<sub>0</sub>-C<sub>4</sub>)alkylenyl; provided that when Ar<sup>3</sup> is phenyl, naphthyl or biphenyl, then R<sup>23</sup> cannot be CONR<sup>25</sup>R<sup>26</sup> where R<sup>25</sup> is hydrogen or Ar<sup>3</sup> and R<sup>26</sup> is Ar<sup>3</sup>;

R<sup>24</sup> is hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl or phenyl optionally independently substituted with up to three hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl or (C<sub>1</sub>-C<sub>4</sub>)alkoxy, wherein said (C<sub>1</sub>-C<sub>4</sub>)alkyl in the definition of R<sup>6</sup> and said (C<sub>1</sub>-C<sub>4</sub>)alkoxy in the definition of R<sup>6</sup> are optionally and independently substituted with up to five fluoro;

Ar<sup>3</sup> is independently defined as set forth for Ar and Ar<sup>1</sup> above;  
said Ar<sup>3</sup> is optionally independently substituted as set forth for Ar and Ar<sup>1</sup> above;

R<sup>27</sup> is hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl;  
R<sup>28</sup> and R<sup>29</sup> are each independently hydrogen, hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro, (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro, phenyl, pyridyl, pyrimidyl, thienyl, furanyl, thiazolyl, oxazolyl, phenoxy, thiophenoxy, SO<sub>2</sub>NR<sup>30</sup>R<sup>31</sup>, CONR<sup>30</sup>R<sup>31</sup> or NR<sup>30</sup>R<sup>31</sup>; said thienyl, pyrimidyl, furanyl, thiazolyl and oxazolyl in the definition of R<sup>28</sup> and R<sup>29</sup> are optionally substituted by up to two hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said phenyl, pyridyl, phenoxy and thiophenoxy in the definition of R<sup>28</sup> and R<sup>29</sup> are optionally substituted by up to three hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-

C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro;

R<sup>30</sup> and R<sup>31</sup> are each independently hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or phenyl, said phenyl is optionally substituted with up to three hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; or

5 R<sup>30</sup> and R<sup>31</sup> are taken together with the nitrogen to which they are attached to form indoliny, pyrrolidiny, piperidiny, piperaziny or morpholiny; said pyrrolidiny and piperidiny in the definition of R<sup>30</sup> and R<sup>31</sup> are optionally substituted with up to two  
10 hydroxy, amino, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said indoliny and piperaziny in the definition of R<sup>30</sup> and R<sup>31</sup> are optionally substituted with up to three hydroxy, amino, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with  
15 up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said morpholiny in the definition of R<sup>30</sup> and R<sup>31</sup> is optionally substituted with up to two substituents independently selected from hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro;

20 A is N optionally substituted with hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl and B is carbonyl; or A is carbonyl and B is N optionally substituted with hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl;

R<sup>32</sup> is hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl;

R<sup>33</sup> is phenyl, pyridyl, pyrimidyl, thiazolyl, oxazolyl, benzyl, quinolyl, isoquinolyl, phthaliziny, quinoxanly, benzothiazoyl, benzoxazolyl, benzofuranyl or benzothienyl;  
25 said phenyl, pyridyl, pyrimidyl, thiazolyl, oxazolyl, benzyl, quinolyl, isoquinolyl, phthaliziny, quinoxanly, benzothiazoyl, benzoxazolyl, benzofuranyl and benzothienyl in the definition of R<sup>33</sup> are optionally substituted with up to three phenyl, phenoxy, NR<sup>34</sup>R<sup>35</sup>, halo, hydroxy, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with  
30 up to five fluoro;

R<sup>34</sup> and R<sup>35</sup> are each independently hydrogen, (C<sub>1</sub>-C<sub>4</sub> alkyl), phenyl or phenylsulfonyl; said phenyl and phenylsulfonyl in the definition of R<sup>34</sup> and R<sup>35</sup> are optionally substituted with up to three halo, hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five  
35 fluoro;

D is CO, CHOH or CH<sub>2</sub>;

E is O, NH or S;

R<sup>36</sup> and R<sup>37</sup> are taken separately and are each independently hydrogen, halo, cyano, hydroxy, amino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino, di-(C<sub>1</sub>-C<sub>6</sub>)alkylamino, pyrrolidino, piperidino, morpholino, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, Ar<sup>4</sup>, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro;

R<sup>38</sup>, R<sup>39</sup> and R<sup>40</sup> are each independently hydrogen or (C<sub>1</sub>-C<sub>4</sub>)-alkyl;

Ar<sup>4</sup> is phenyl, furanyl, thienyl, pyridyl, pyrimidyl, pyrazinyl or pyridazinyl; said Ar<sup>4</sup>

being optionally substituted with up to three hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; or

R<sup>36</sup> and R<sup>37</sup> are taken together on adjacent carbon atoms and are -O-(CH<sub>2</sub>)<sub>t</sub>-O-; t is 1, 2 or 3;

Y is (C<sub>2</sub>-C<sub>6</sub>)alkylene;

R<sup>44</sup>, R<sup>45</sup> and R<sup>46</sup> are each independently hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl;

m and n are each independently 1, 2 or 3, provided that the sum of m and n is 2, 3 or 4;

k is 0, 1, 2, 3 or 4;

Y<sup>1</sup> is a covalent bond, carbonyl, sulfonyl or oxycarbonyl;

R<sup>43</sup> is (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, Ar<sup>5</sup>-(C<sub>0</sub>-C<sub>4</sub>)alkylenyl, NR<sup>47</sup>R<sup>48</sup> or (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted with one to five fluoro; provided that when Y<sup>1</sup> is a covalent bond or oxycarbonyl, then R<sup>43</sup> is not NR<sup>47</sup>R<sup>48</sup>;

R<sup>47</sup> and R<sup>48</sup> are taken separately and are each independently selected from

hydrogen, Ar<sup>5</sup>, (C<sub>1</sub>-C<sub>6</sub>)alkyl and Ar<sup>5</sup>-(C<sub>0</sub>-C<sub>4</sub>)alkylenyl; or

R<sup>47</sup> and R<sup>48</sup> are taken together with the nitrogen atom to which they are attached to form azetidiny, pyrrolidiny, piperidiny, piperaziny, morpholiny, azepiny, azabicyclo[3.2.2]nonany, azabicyclo[2.2.1]heptyl, 1,2,3,4-tetrahydroisoquinoly, 6,7-dihydro-5H-dibenzo[c,e]azepiny or 5,6,7,8-tetrahydropyrido[4,3-d]pyrimidy; said

azetidiny in the definition of R<sup>47</sup> and R<sup>48</sup> are optionally substituted with one hydroxy, amino, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said pyrrolidiny, piperidiny and azepiny in the definition of R<sup>47</sup> and R<sup>48</sup> are optionally substituted with up to two hydroxy, amino, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-


C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-



- C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said morpholinyl in the definition of R<sup>47</sup> and R<sup>48</sup> is optionally substituted with up to two substituents independently selected from hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said piperazinyl, 1,2,3,4-tetrahydroisoquinolyl and 5,6,7,8-tetrahydro[4,3-d]pyrimidyl in the definition of R<sup>47</sup> and R<sup>48</sup> are optionally substituted with up to three hydroxy, amino, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; and said 6,7-dihydro-5H-dibenzo[c,e]azepinyl in the definition of R<sup>47</sup> and R<sup>48</sup> are optionally substituted with up to four hydroxy, amino, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro;
- Ar<sup>5</sup> is independently defined as set forth for Ar and Ar<sup>1</sup> above;
- Ar<sup>5</sup> is optionally independently substituted as set forth for Ar and Ar<sup>1</sup> above;
- R<sup>42</sup> and R<sup>42a</sup> are independently hydrogen, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, Ar<sup>6</sup>-(C<sub>0</sub>-C<sub>3</sub>)alkylenyl, Ar<sup>6</sup>-(C<sub>2</sub>-C<sub>4</sub>)alkenyl, Ar<sup>6</sup>-carbonyl or (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted with up to five fluoro;
- Ar<sup>6</sup> is independently defined as set forth for Ar and Ar<sup>1</sup> above;
- Ar<sup>6</sup> is optionally independently substituted as set forth for Ar and Ar<sup>1</sup> above; and
- R<sup>41</sup> and R<sup>41a</sup> are each independently hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl.

2. A compound of claim 1, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, wherein:

R<sup>3</sup> is



substituted by R<sup>18</sup>, R<sup>19</sup> or R<sup>20</sup>;


G<sup>3</sup>, G<sup>4</sup> and G<sup>5</sup> are taken separately and are each hydrogen, r is 0 and R<sup>18</sup> is hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl or phenyl optionally substituted by up to three hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl

- optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; R<sup>19</sup> and R<sup>20</sup> are each independently (C<sub>1</sub>-C<sub>4</sub>)alkyl;
- G<sup>3</sup>, G<sup>4</sup> and G<sup>5</sup> are taken separately and are each hydrogen; r is 1; and R<sup>18</sup> is hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl or phenyl optionally substituted by up
- 5 to three hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; R<sup>19</sup> and R<sup>20</sup> are each independently hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl; or G<sup>3</sup> and G<sup>4</sup> are taken together and are (C<sub>1</sub>-C<sub>3</sub>)alkylene; r is 0 or 1; and R<sup>18</sup>, R<sup>19</sup>, R<sup>20</sup> and G<sup>5</sup> are hydrogen; or
- 10 G<sup>4</sup> and G<sup>5</sup> are taken together and are (C<sub>1</sub>-C<sub>3</sub>)alkylene; r is 0 or 1; and R<sup>18</sup>, R<sup>19</sup>, R<sup>20</sup> and G<sup>3</sup> are hydrogen;
- R<sup>17</sup> is SO<sub>2</sub>NR<sup>21</sup>R<sup>22</sup>, CONR<sup>21</sup>R<sup>22</sup>, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl, Ar<sup>2</sup>-carbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfinyl, Ar<sup>2</sup>-sulfonyl, Ar<sup>2</sup>-sufinyl and (C<sub>1</sub>-C<sub>6</sub>)alkyl;
- 15 R<sup>21</sup> and R<sup>22</sup> are taken separately and are each independently selected from hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl and Ar<sup>2</sup>-(C<sub>0</sub>-C<sub>4</sub>)alkylenyl; or R<sup>21</sup> and R<sup>22</sup> are taken together with the nitrogen atom to which they are attached to form azetidiny, pyrrolidiny, piperidiny, piperaziny, morpholiny, azepiny, azabicyclo[3.2.2]nonany, azabicyclo[2.2.1]heptyl, 6,7-dihydro-5H-
- 20 dibenzo[c,e]azepiny, 1,2,3,4-tetrahydro-isoquinoly or 5,6,7,8-tetrahydropyrido[4,3-d]pyrimidy; said azetidiny in the definition of R<sup>21</sup> and R<sup>22</sup> is optionally substituted independently with one substituent selected from hydroxy, amino, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said pyrrolidiny, piperidiny, morpholiny, azepiny in the definition of R<sup>21</sup> and R<sup>22</sup> are optionally
- 25 substituted independently with up to two substituents independently selected from hydroxy, amino, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said morpholiny in the definition of R<sup>21</sup> and R<sup>22</sup> is optionally
- 30 substituted with up to two substituents independently selected from hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said piperaziny in the definition of R<sup>21</sup> and R<sup>22</sup> is optionally substituted independently with up to three substituents independently selected from phenyl, pyridyl, pyrimidy, (C<sub>1</sub>-
- 35 C<sub>4</sub>)alkoxycarbonyl and (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro; said

1,2,3,4-tetrahydro-isoquinolyl and said 5,6,7,8-tetrahydropyrido[4,3-d]pyrimidyl in the definition of  $R^{21}$  and  $R^{22}$  are optionally substituted independently with up to three substituents independently selected from hydroxy, amino, halo, hydroxy-( $C_1$ - $C_4$ )alkyl, ( $C_1$ - $C_4$ )alkoxy-( $C_1$ - $C_4$ )alkyl, ( $C_1$ - $C_4$ )alkyl optionally substituted with up to five fluoro and ( $C_1$ - $C_4$ )alkoxy optionally substituted with up to five fluoro; said pyrimidyl, pyridyl and phenyl which are optionally substituted on said piperazine in the definition of  $R^{21}$  and  $R^{22}$  is optionally substituted with up to three substituents selected from hydroxy, amino, hydroxy-( $C_1$ - $C_4$ )alkyl, ( $C_1$ - $C_4$ )alkoxy-( $C_1$ - $C_4$ )alkyl, ( $C_1$ - $C_4$ )alkyl optionally substituted with up to five fluoro and ( $C_1$ - $C_4$ )alkoxy optionally substituted with up to five fluoro; and said 6,7-dihydro-5H-dibenzo[c,e]azepinyl in the definition of  $R^{21}$  and  $R^{22}$  is optionally substituted with up to four substituents independently selected from hydroxy, amino, halo, hydroxy-( $C_1$ - $C_4$ )alkyl, ( $C_1$ - $C_4$ )alkoxy-( $C_1$ - $C_4$ )alkyl, ( $C_1$ - $C_4$ )alkyl optionally substituted with up to five fluoro and ( $C_1$ - $C_4$ )alkoxy optionally substituted with up to five fluoro.

3. A compound of claim 1, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, wherein:

$R^3$  is




$R^{23}$  is  $CONR^{25}R^{26}$ ,  $SO_2R^{25}R^{26}$ , wherein  $R^{25}$  is hydrogen ( $C_1$ - $C_4$ )alkyl or  $Ar^3$ -( $C_0$ - $C_4$ )alkylenyl and  $R^{26}$  is  $Ar^3$ -( $C_0$ - $C_4$ )alkylenyl; provided that when  $Ar^3$  is phenyl, naphthyl or biphenyl, then  $R^{23}$  cannot be  $CONR^{25}R^{26}$  where  $R^{25}$  is hydrogen or  $Ar^3$  and  $R^{26}$  is  $Ar^3$ ;

$R^{24}$  is hydrogen, ( $C_1$ - $C_4$ )alkyl, ( $C_1$ - $C_4$ )alkoxycarbonyl or phenyl optionally substituted by up to three ( $C_1$ - $C_4$ )alkyl optionally substituted with up to five fluoro, ( $C_1$ - $C_4$ )alkoxy optionally substituted with up to five fluoro, hydroxy, halo or hydroxy-( $C_1$ - $C_3$ )alkyl.

4. A compound of claim 1, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, wherein:

$R^3$  is



$R^{27}$  is hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl;

$R^{28}$  and  $R^{29}$  are each independently hydrogen, hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro,

5 (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro, phenyl, pyridyl, pyrimidyl, thienyl, furanyl, thiazolyl, oxazolyl, phenoxy, thiophenoxy, SO<sub>2</sub>NR<sup>30</sup>R<sup>31</sup>, CONR<sup>30</sup>R<sup>31</sup> or NR<sup>30</sup>R<sup>31</sup>; said thienyl, pyrimidyl, furanyl, thiazolyl and oxazolyl in the definition of  $R^{28}$  and  $R^{29}$  are optionally substituted by up to two hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl,

10 (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said phenyl, pyridyl, phenoxy and thiophenoxy in the definition of  $R^{28}$  and  $R^{29}$  are optionally substituted by up to three hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro;

15  $R^{30}$  and  $R^{31}$  are each independently hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or phenyl, said phenyl is optionally substituted with up to three hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; or


20  $R^{30}$  and  $R^{31}$  are taken together with the nitrogen to which they are attached to form indoliny, pyrrolidiny, piperidiny, piperaziny or morpholiny; said pyrrolidiny and piperidiny in the definition of  $R^{30}$  and  $R^{31}$  are optionally substituted with up to two hydroxy, amino, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said indoliny and piperaziny in the definition of  $R^{30}$  and  $R^{31}$  are

25 optionally substituted with up to three hydroxy, amino, or hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; and said morpholiny in the definition of  $R^{30}$  and  $R^{31}$  is optionally substituted with up to two substituents independently selected from hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-

C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro.

5. A compound of claim 1, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, wherein:

5 R<sup>3</sup> is



A is N optionally substituted with hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl and B is carbonyl; or

A is carbonyl and B is N optionally substituted with hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl;

R<sup>32</sup> is hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl;


10 R<sup>33</sup> is phenyl, pyridyl, pyrimidyl, thiazolyl, oxazolyl, benzyl, quinolyl, isoquinolyl, phthaliziny, quinoxanly, benzothiazoyl, benzoxazolyl, benzofuranyl or benzothieryl; said phenyl, pyridyl, pyrimidyl, thiazolyl, oxazolyl, benzyl, quinolyl, isoquinolyl, phthaliziny, quinoxanly, benzothiazoyl, benzoxazolyl, benzofuranyl and benzothieryl in the definition of R<sup>33</sup> are optionally substituted with up to three phenyl, phenoxy, NR<sup>34</sup>R<sup>35</sup>, halo, hydroxy, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro;

R<sup>34</sup> and R<sup>35</sup> are each independently hydrogen, (C<sub>1</sub>-C<sub>4</sub> alkyl), phenyl or phenylsulfonyl; said phenyl and phenylsulfonyl in the definition of R<sup>34</sup> and R<sup>35</sup> are


20 optionally substituted with up to three halo, hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro.

6. A compound of claim 1, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, wherein:

25 R<sup>3</sup> is



or



D is CO, CHOH or CH<sub>2</sub>;

E is O, NH or S;

R<sup>36</sup> and R<sup>37</sup> are taken separately and are each independently hydrogen, halo, cyano, hydroxy, amino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino, di-(C<sub>1</sub>-C<sub>6</sub>)alkylamino, pyrrolidino, piperidino, morpholino, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, Ar<sup>4</sup>, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro;

R<sup>38</sup>, R<sup>39</sup> and R<sup>40</sup> are each independently hydrogen or (C<sub>1</sub>-C<sub>4</sub>)-alkyl;


Ar<sup>4</sup> is phenyl, furanyl, thienyl, pyridyl, pyrimidyl, pyrazinyl or pyridazinyl; said Ar<sup>4</sup> being optionally substituted with up to three hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; or

R<sup>36</sup> and R<sup>37</sup> are taken together on adjacent carbon atoms and are -O-(CH<sub>2</sub>)<sub>t</sub>-O-;

t is 1, 2 or 3.

7. A compound of claim 1, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, wherein:

R<sup>3</sup> is



Y is (C<sub>2</sub>-C<sub>6</sub>)alkylene;

R<sup>44</sup>, R<sup>45</sup> and R<sup>46</sup> are each independently hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl;

m and n are each independently 1, 2 or 3, provided that the sum of m and n is 2, 3 or

4;

k is 0 to 4;

Y<sup>1</sup> is a covalent bond, carbonyl, sulfonyl or oxycarbonyl;

R<sup>43</sup> is (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, Ar<sup>5</sup>-(C<sub>0</sub>-C<sub>4</sub>)alkylenyl, NR<sup>47</sup>R<sup>48</sup> or (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted with one to five fluoro; provided that when Y<sup>1</sup> is a covalent bond or

10 oxycarbonyl, then R<sup>43</sup> is not NR<sup>47</sup>R<sup>48</sup>;

R<sup>47</sup> and R<sup>48</sup> are taken separately and are each independently selected from hydrogen, Ar<sup>5</sup>, (C<sub>1</sub>-C<sub>6</sub>)alkyl and Ar<sup>5</sup>-(C<sub>0</sub>-C<sub>4</sub>)alkylenyl; or

R<sup>47</sup> and R<sup>48</sup> are taken together with the nitrogen atom to which they are attached to form azetidiny, pyrrolidiny, piperidiny, piperaziny, morpholiny, azepiny,

15 azabicyclo[3.2.2]nonany, azabicyclo[2.2.1]heptyl, 1,2,3,4-tetrahydroisoquinoly, 6,7-dihydro-5H-dibenzo[c,e]azepiny or 5,6,7,8-tetrahydropyrido[4,3-d]pyrimidy; said

azetidiny in the definition of R<sup>47</sup> and R<sup>48</sup> are optionally substituted with one hydroxy, amino, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five


fluoro; said pyrrolidinyl, piperidinyl and azepinyl in the definition of  $R^{47}$  and  $R^{48}$  are optionally substituted with up to two hydroxy, amino, hydroxy-( $C_1$ - $C_4$ )alkyl, ( $C_1$ - $C_4$ )alkoxy-( $C_1$ - $C_4$ )alkyl, ( $C_1$ - $C_4$ )alkyl optionally substituted with up to five fluoro or ( $C_1$ - $C_4$ )alkoxy optionally substituted with up to five fluoro; said morpholinyl in the  
5 definition of  $R^{47}$  and  $R^{48}$  is optionally substituted with up to two substituents independently selected from hydroxy-( $C_1$ - $C_4$ )alkyl, ( $C_1$ - $C_4$ )alkoxy-( $C_1$ - $C_4$ )alkyl, ( $C_1$ - $C_4$ )alkyl optionally substituted with up to five fluoro and ( $C_1$ - $C_4$ )alkoxy optionally substituted with up to five fluoro; said piperazinyl, 1,2,3,4-tetrahydroisoquinolyl and 5,6,7,8-tetrahydro[4,3-d]pyrimidyl in the definition of  $R^{47}$  and  $R^{48}$  are optionally  
10 substituted with up to three hydroxy, amino, halo, hydroxy-( $C_1$ - $C_4$ )alkyl, ( $C_1$ - $C_4$ )alkoxy-( $C_1$ - $C_4$ )alkyl, ( $C_1$ - $C_4$ )alkyl optionally substituted with up to five fluoro or ( $C_1$ - $C_4$ )alkoxy optionally substituted with up to five fluoro; and said 6,7-dihydro-5H-dibenzo[c,e]azepinyl in the definition of  $R^{47}$  and  $R^{48}$  are optionally substituted with up to four hydroxy, amino, halo, hydroxy-( $C_1$ - $C_4$ )alkyl, ( $C_1$ - $C_4$ )alkoxy-( $C_1$ - $C_4$ )alkyl, ( $C_1$ - $C_4$ )alkyl optionally substituted with up to five fluoro or ( $C_1$ - $C_4$ )alkoxy optionally  
15 substituted with up to five fluoro.

8. A compound of claim 7, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein:

$R^1$  is (R)-1-hydroxy-ethyl;

20  $R^2$  is hydrogen;

$R^3$  is



k is 0;

$Y^1$  is a covalent bond; and


25  $R^{43}$  is 4-pyrimidinyl substituted at the 2-position with 1-hydroxymethyl.

9. The compound of claim 8, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, which is 1R-(4-{1'-[2-(1R-Hydroxy-ethyl)-pyrimidin-4-yl]-[4,4']bipiperidin-1-yl}-pyrimidin-2-yl)-ethanol.




10. A compound of claim 1, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, wherein:

R<sup>3</sup> is



or



5 R<sup>42</sup> and R<sup>42a</sup> are independently hydrogen, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, Ar<sup>6</sup>-(C<sub>0</sub>-C<sub>3</sub>)alkylenyl, Ar<sup>6</sup>-(C<sub>2</sub>-C<sub>4</sub>)alkenyl, Ar<sup>6</sup>-carbonyl or (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted with up to five fluoro; and


R<sup>41</sup> and R<sup>41a</sup> are independently is hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl.

11. A compound of claim 1, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, wherein:

R<sup>1</sup> is C(OH)R<sup>4</sup>R<sup>5</sup>, where R<sup>4</sup> and R<sup>5</sup> are each independently hydrogen or methyl;

R<sup>2</sup> is hydrogen;

R<sup>3</sup> is



15 wherein said R<sup>3</sup> is substituted by R<sup>6</sup>, R<sup>7</sup> or R<sup>8</sup>;

G, G<sup>1</sup> and G<sup>2</sup> are taken separately and are each hydrogen and R<sup>6</sup> is hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl; R<sup>7</sup> and R<sup>8</sup> are each independently hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl; or G and G<sup>1</sup> are taken together and are (C<sub>1</sub>-C<sub>3</sub>)alkylene and R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup> and G<sup>2</sup> are hydrogen; or

20 G<sup>1</sup> and G<sup>2</sup> are taken together and are (C<sub>1</sub>-C<sub>3</sub>)alkylene and R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup> and G are hydrogen;

q is 0 or 1;

X is a covalent bond, oxycarbonyl, vinylenylcarbonyl, oxy(C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl, thio(C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl or vinylenylsulfonyl; said vinylenylcarbonyl and said

25 vinylenylsulfonyl in the definition of X are optionally substituted on one or two

vinylene carbons with (C<sub>1</sub>-C<sub>4</sub>)alkyl, benzyl or Ar; said oxy(C<sub>1</sub>-C<sub>4</sub>)alkylene carbonyl and said thio(C<sub>1</sub>-C<sub>4</sub>)alkylene carbonyl in the definition of X are optionally substituted with up to two (C<sub>1</sub>-C<sub>4</sub>)alkyl, benzyl or Ar;

R<sup>9</sup> is (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, Ar<sup>1</sup>-(C<sub>0</sub>-C<sub>4</sub>)alkylene or (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted with

5 up to five fluoro;

Ar<sup>1</sup> is phenyl, naphthyl, pyridyl, pyrimidyl, pyrazinyl, pyridazinyl, triazinyl, quinolyl, isoquinolyl, quinazolyl, quinoxalyl, phthalazinyl, cinnolyl, naphthyridinyl, pteridinyl, pyrazinopyrazinyl, pyrazinopyridazinyl, pyrimidopyridazinyl, pyrimidopyrimidyl, pyridopyrimidyl, pyridopyrazinyl, pyridopyridazinyl, pyrrolyl, furanyl, thienyl, imidazolyl, oxazolyl, thiazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzoxazolyl, benzothiazolyl, indazolyl, benzisoxazolyl, benzisothiazolyl, pyrrolopyridyl, furopyridyl, thienopyridyl, imidazolopyridyl, oxazolopyridyl, thiazolopyridyl, pyrazolopyridyl, isoxazolopyridyl, isothiazolopyridyl, pyrrolopyrimidyl, furopyrimidyl, thienopyrimidyl, imidazolopyrimidyl, oxazolopyrimidyl, thiazolopyrimidyl, pyrazolopyrimidyl, isoxazolopyrimidyl, isothiazolopyrimidyl, pyrrolopyrazinyl, furopyrazinyl, thienopyrazinyl, imidazolopyrazinyl, oxazolopyrazinyl, thiazolopyrazinyl, pyrazolopyrazinyl, isoxazolopyrazinyl, isothiazolopyrazinyl, pyrrolopyridazinyl, furopyridazinyl, thienopyridazinyl, imidazolopyridazinyl, oxazolopyridazinyl, thiazolopyridazinyl, pyrazolopyridazinyl, isoxazolopyridazinyl or isothiazolopyridazinyl; and said Ar<sup>1</sup> is optionally substituted as set forth in claim 1.

12. A compound of claim 11, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, wherein:

25 X is a covalent bond, oxycarbonyl or vinylenyl carbonyl optionally substituted on one or two vinylenyl carbons with (C<sub>1</sub>-C<sub>4</sub>)alkyl, benzyl or Ar;

R<sup>9</sup> is Ar<sup>1</sup>-(C<sub>0</sub>-C<sub>4</sub>)alkylene;

Ar<sup>1</sup> is phenyl, naphthyl, pyridyl, pyrimidyl, pyrazinyl, triazinyl, quinolyl, isoquinolyl, quinazolyl, quinoxalyl, furanyl, thienyl, indolyl, benzofuranyl, benzothienyl, benzoxazolyl, benzothiazolyl, furopyridyl, oxazolopyridyl, thiazolopyridyl, thienopyridyl, furopyrimidyl, thienopyrimidyl, oxazolopyrimidyl or thiazolopyrimidyl; and

said Ar<sup>1</sup> is optionally substituted as set forth in claim 1.

13. A compound of claim 12, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, wherein:

R<sup>2</sup> is hydrogen;

R<sup>4</sup> is hydrogen or methyl;

R<sup>5</sup> is methyl;

G, G<sup>1</sup> and G<sup>2</sup> are hydrogen;


5 R<sup>6</sup> and R<sup>7</sup> are each independently hydrogen or methyl;

R<sup>8</sup> is hydrogen.

14. A compound of claim 13, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein:

R<sup>1</sup> is (R)-1-hydroxy-ethyl; and

10 R<sup>3</sup> is



15. The compound of claim 14, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, wherein: R<sup>9</sup> is 2-furo[3,2-c]pyridyl.

16. The compound of claim 14, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 2-(4-chloro-furo[3,2-c]pyridyl).

17. The compound of claim 14, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 2-(4-pyrrolidin-1-yl-furo[3,2-c]pyridyl).

20 18. The compound of claim 14, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 2-(4-morpholin-4-yl-furo[3,2-c]pyridyl).

19. The compound of claim 14, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 2-imidazo[1,2-a]pyridyl.


20. A compound of claim 14, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, selected from furo[3,2-c]pyridin-2-yl-{4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-3R,5S-dimethyl-piperazin-1-yl}-methanone; (4-chloro-furo[3,2-c]pyridin-2-yl)-{4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-3R,5S-dimethyl-piperazin-1-yl}-methanone; {4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-3R,5S-

dimethyl-piperazin-1-yl)-(4-pyrrolidin-1-yl-furo[3,2-c]pyridin-2-yl)-methanone; {4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-3R,5S-dimethyl-piperazin-1-yl)-(4-morpholin-4-yl-furo[3,2-c]pyridin-2-yl)-methanone; and {4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-3R,5S-dimethyl-piperazin-1-yl}-imidazo[1,2-a]pyridin-2-yl-methanone.

- 5            21.     A compound of claim 13, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein:

$R^1$  is (R)-1-hydroxy-ethyl; and

$R^3$  is




- 10           22.     The compound of claim 21, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein:  $R^9$  is 2-furo[3,2-c]pyridyl.

23.     The compound of claim 21, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, which is furo[3,2-c]pyridin-2-yl-{4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl}-methanone.

- 15           24.     A compound of claim 13, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein:

$R^1$  is (R)-1-hydroxy-ethyl; and

$R^3$  is



- 20           25.     The compound of claim 24, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein:  $R^9$  is 3-pyridyl.

26.     The compound of claim 24, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein:  $R^9$  is 3-(2-methylpyridyl).

- 25           27.     The compound of claim 24, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein:  $R^9$  is 3-(5-chloropyridyl).


28. The compound of claim 24, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein:  $R^9$  is 3-(6-methylpyridyl).

29. A compound of claim 24, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, selected from 4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazine-1-carboxylic acid pyridin-3-yl ester; 4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazine-1-carboxylic acid 2-methyl-pyridin-3-yl ester; 4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazine-1-carboxylic acid 5-chloro-pyridin-3-yl ester; and 4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazine-1-carboxylic acid 6-methyl-pyridin-3-yl ester.

30. A compound of claim 13, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein:

$R^1$  is (R)-1-hydroxy-ethyl; and

$R^3$  is




31. The compound of compound of claim 30, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein:  $R^9$  is 2-thienyl.

32. The compound of claim 30, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, which is (E)-1-{4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl}-3-thiophen-2-yl-propenone.

33. A compound of claim 13, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein:

$R^1$  is (R)-1-hydroxy-ethyl;

$R^3$  is



; and

R<sup>9</sup> is pyrimidyl or triazinyl; said pyrimidyl or triazinyl is optionally substituted with up to two hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkylenyl, phenyl, piperazinyl optionally substituted with (C<sub>1</sub>-C<sub>4</sub>)alkyl, or imidazolyl optionally substituted with up to two (C<sub>1</sub>-C<sub>4</sub>)alkyl.

5            34.      A compound of claim 33, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is pyrimid-2-yl optionally substituted with up to two (C<sub>1</sub>-C<sub>4</sub>)alkyl, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl or (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl.

10           35.      The compound of claim 34, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4,6-dimethylpyrimid-2-yl.

36.      The compound of claim 34, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4-methoxymethyl-6-methylpyrimid-2-yl.

15           37.      The compound of claim 34, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4-hydroxymethyl-6-methylpyrimid-2-yl.

20           38.      A compound of claim 34, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, selected from 1R-{4-[4-(4,6-dimethyl-pyrimidin-2-yl)-3R,5S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol; 1R-{4-[4-(4-methoxymethyl-6-methyl-pyrimidin-2-yl)-3R,5S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol; and 1R-{4-[4-(4-hydroxymethyl-6-methyl-pyrimidin-2-yl)-3R,5S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol.

25           39.      A compound of claim 33, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is pyrimid-4-yl optionally substituted with up to two (C<sub>1</sub>-C<sub>4</sub>)alkylpiperazin-1-yl or imidazolyl; and said imidazolyl is optionally substituted with up to two (C<sub>1</sub>-C<sub>4</sub>)alkyl.

30           40.      The compound of claim 39, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 2-(4-methylpiperazin-1-yl)-pyrimid-4-yl.

41.      The compound of claim 39, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 2-(4-ethylpiperazin-1-yl)-pyrimid-4-yl.

42. The compound of claim 39, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 2-(4-methylimidazol-1-yl)-pyrimid-4-yl.

43. The compound of claim 39, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 2-(2-methylimidazol-1-yl)-pyrimid-4-yl.

44. The compound of claim 39, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 2-(2,4-dimethylimidazol-1-yl)-pyrimid-4-yl.

45. The compound of claim 39, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 2-(4-isopropylpiperazin-1-yl)-pyrimid-4-yl.

46. A compound of claim 39, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, selected from 1R-(4-{3R,5S-dimethyl-4-[2-(4-methyl-piperazin-1-yl)-pyrimidin-4-yl]-piperazin-1-yl}-pyrimidin-2-yl)-ethanol; 1R-(4-{4-[2-(4-ethyl-piperazin-1-yl)-pyrimidin-4-yl]-3R,5S-dimethyl-piperazin-1-yl}-pyrimidin-2-yl)-ethanol; 1R-(4-{3R,5S-dimethyl-4-[2-(4-methyl-imidazol-1-yl)-pyrimidin-4-yl]-piperazin-1-yl}-pyrimidin-2-yl)-ethanol; 1R-(4-{3R,5S-dimethyl-4-[2-(2-methyl-imidazol-1-yl)-pyrimidin-4-yl]-piperazin-1-yl}-pyrimidin-2-yl)-ethanol; 1R-(4-{4-[2-(2,4-dimethyl-imidazol-1-yl)-pyrimidin-4-yl]-3R,5S-dimethyl-piperazin-1-yl}-pyrimidin-2-yl)-ethanol; and 1R-(4-{4-[2-(4-isopropyl-piperazin-1-yl)-pyrimidin-4-yl]-3R,5S-dimethyl-piperazin-1-yl}-pyrimidin-2-yl)-ethanol.

47. A compound of claim 33, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is [1,3,5]-triazin-2-yl optionally substituted with up to two (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkylpiperazin-1-yl or phenyl.

48. The compound of claim 47, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4-methyl-6-(4-methylpiperazin-1-yl)-[1,3,5]-triazin-2-yl.

49. The compound of claim 47, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4-methoxy-6-methyl-[1,3,5]-triazin-2-yl.

50. The compound of claim 47, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4,6-dimethoxy-[1,3,5]-triazin-2-yl.

51. The compound of claim 47, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4-ethoxy-6-methyl-[1,3,5]-triazin-2-yl.

52. The compound of claim 47, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4-isopropoxy-6-methyl-[1,3,5]-triazin-2-yl.

53. The compound of claim 47, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4-phenyl-[1,3,5]-triazin-2-yl.

54. The compound of claim 47, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4-hydroxymethyl-6-methoxy-[1,3,5]-triazin-2-yl.

55. The compound of claim 47, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4-isopropoxy-6-methoxy-[1,3,5]-triazin-2-yl.

56. The compound of claim 47, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4-isopropyl-[1,3,5]-triazin-2-yl.

57. The compound of claim 47, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4-ethyl-6-methoxy-[1,3,5]-triazin-2-yl.

58. The compound of claim 47, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4-cyclopropyl-[1,3,5]-triazin-2-yl.

59. The compound of claim 47, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4,6-dimethyl-[1,3,5]-triazin-2-yl.

60. The compound of claim 47, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4-methyl-6-phenyl-[1,3,5]-triazin-2-yl.

61. A compound of claim 47, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, selected from 1R-(4-{3R,5S-dimethyl-4-[4-methyl-6-(4-methyl-piperazin-1-yl)-[1,3,5]triazin-2-yl]-piperazin-1-yl}-pyrimidin-2-yl)-ethanol; 1R-{4-[4-(4-methoxy-6-methyl-[1,3,5]triazin-2-yl)-3R,5S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol; 1R-{4-[4-(4,6-dimethoxy-




[1,3,5]triazin-2-yl)-3R,5S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol; 1R-{4-[4-(4-ethoxy-6-methyl-[1,3,5]triazin-2-yl)-3R,5S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol; 1R-{4-[4-(4-isopropoxy-6-methyl-[1,3,5]triazin-2-yl)-3R,5S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol; 1R-{4-[3R,5S-dimethyl-4-(4-phenyl-[1,3,5]triazin-2-yl)-piperazin-1-yl]-pyrimidin-2-yl}-ethanol; 1R-{4-[4-(4-hydroxymethyl-6-methoxy-[1,3,5]triazin-2-yl)-3R,5S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol; 1R-{4-[4-(4-isopropoxy-6-methoxy-[1,3,5]triazin-2-yl)-3R,5S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol; 1R-{4-[4-(4-isopropyl-[1,3,5]triazin-2-yl)-3R,5S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol; 1R-{4-[4-(4,6-dimethyl-[1,3,5]triazin-2-yl)-3R,5S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol; 1R-{4-[3R,5S-dimethyl-4-(4-methyl-6-phenyl-[1,3,5]triazin-2-yl)-piperazin-1-yl]pyrimidin-2-yl}-ethanol; 1R-{4-[4-(4-cyclopropyl-[1,3,5]triazin-2-yl)-3R,5S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol; and 1R-{4-[4-(4-ethyl-6-methoxy-[1,3,5]triazin-2-yl)-3R,5S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol.

62. A compound of claim 13, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein:

R<sup>1</sup> is (R)-1-hydroxy-ethyl;

R<sup>3</sup> is



; and

R<sup>9</sup> is pyrimidyl or triazinyl, said pyrimidyl and triazinyl optionally substituted with up to two hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, triazolyl, acetyl, morpholinyl, (C<sub>1</sub>-C<sub>4</sub>)alkylpiperazinyl, phenyl or imidazolyl optionally substituted with up to two (C<sub>1</sub>-C<sub>4</sub>)alkyl.

63. A compound of claim 62, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is pyrimid-2-yl optionally substituted with up to two (C<sub>1</sub>-C<sub>4</sub>)alkyl, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl or triazolyl.

64. The compound of claim 63, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4,6-dimethyl-pyrimid-2-yl.

65. The compound of claim 63, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4-hydroxymethyl-6-methylpyrimid-2-yl.

66. The compound of claim 63, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4-[1,2,4]-triazol-1-yl-pyrimid-2-yl.

67. A compound of claim 63, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, selected from 1R-{4-[4-(4,6-dimethyl-pyrimidin-2-yl)-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol; 1R-{4-[4-(4-hydroxymethyl-6-methyl-pyrimidin-2-yl)-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol; and 1R-{4-[2R,6S-dimethyl-4-(4-[1,2,4]triazol-1-yl-pyrimidin-2-yl)-piperazin-1-yl]-pyrimidin-2-yl}-ethanol.

68. A compound of claim 62, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is pyrimid-4-yl optionally substituted with up to two (C<sub>1</sub>-C<sub>4</sub>)alkyl, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, acetyl, morpholinyl, (C<sub>1</sub>-C<sub>4</sub>)alkylpiperazinyl, triazolyl or imidazolyl optionally substituted with up to two (C<sub>1</sub>-C<sub>4</sub>)alkyl.

69. The compound of claim 68, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 2,6-dimethyl-pyrimid-4-yl.

70. The compound of claim 68, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 2-hydroxymethyl-6-methyl-pyrimid-4-yl.

71. The compound of claim 68, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 2-acetyl-pyrimid-4-yl.

72. The compound of claim 68, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 2-morpholin-4-yl-pyrimid-4-yl.

73. The compound of claim 68, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 2-(4-methylpiperazin-1-yl)-pyrimid-4-yl.

74. The compound of claim 68, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 2-[1,2,4]-triazol-1-yl-pyrimid-4-yl.

75. The compound of claim 68, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 2-(1S-hydroxyethyl)-pyrimid-4-yl.

76. The compound of claim 68, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 2-(1R-hydroxyethyl)-pyrimid-4-yl.

77. The compound of claim 68, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 2-(4-ethylpiperazin-1-yl)-pyrimid-4-yl.


78. The compound of claim 68, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 2-(4-methylimidazol-1-yl)-pyrimid-4-yl.

79. The compound of claim 68, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 2-(2,4-dimethylimidazol-1-yl)-pyrimid-4-yl.

80. A compound of claim 13, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, selected from 1R-{4-[4-(2,6-dimethyl-pyrimidin-4-yl)-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol; 1R-(4-{4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl}-pyrimidin-2-yl)-ethanol; 1R-(4-{4-[2-(1S-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl}-pyrimidin-2-yl)-ethanol; 1S-(4-{4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl}-pyrimidin-2-yl)-ethanol; 1-{4-[4-(2-acetyl-pyrimidin-4-yl)-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanone; 1RS-(4-{4-[2-(1RS-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl}-pyrimidin-2-yl)-ethanol; (4-{4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-3R,5S-dimethyl-piperazin-1-yl}-pyrimidin-2-yl)-ethanone; 1R-(4-[2R,6S-dimethyl-4-(2-morpholin-4-yl)-pyrimidin-4-yl]-piperazin-1-yl)-pyrimidin-2-yl)-ethanol; 1R-(4-{2R,6S-dimethyl-4-[2-(4-methyl-piperazin-1-yl)-pyrimidin-4-yl]-piperazin-1-yl}-pyrimidin-2-yl)-ethanol; 1R-(4-[2R,6S-dimethyl-4-(2-[1,2,4]triazol-1-yl)-pyrimidin-4-yl]-piperazin-1-yl)-pyrimidin-2-yl)-ethanol; 1R-(4-{4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6R-dimethyl-piperazin-1-yl}-pyrimidin-2-yl)-ethanol; 1R-(4-{4-[2-(4-ethyl-piperazin-1-yl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl}-pyrimidin-2-yl)-ethanol; 1R-(4-{2R,6S-dimethyl-4-[2-(4-methyl-imidazol-1-yl)-pyrimidin-4-yl]-piperazin-1-yl}-pyrimidin-2-yl)-ethanol; 1R-(4-[4-(4-cyclopropyl-[1,3,5]triazin-2-yl)-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl)-

ethanol; 1R-{4-[4-(4,6-dimethyl-[1,3,5]triazin-2-yl)-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol; 1R-{4-[4-(4-hydroxymethyl-6-phenyl-[1,3,5]triazin-2-yl)-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol; 1R-{4-[4-(4-methoxy-6-methoxymethyl-[1,3,5]triazin-2-yl)-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol; 1R-{4-[2R,6S-dimethyl-4-(4-methyl-[1,3,5]triazin-2-yl)-piperazin-1-yl]-pyrimidin-2-yl}-ethanol; and 1R-(4-{4-[2-(2,4-dimethyl-imidazol-1-yl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl}-pyrimidin-2-yl)-ethanol.

81. A compound of claim 13 wherein R<sup>1</sup> is (R)-1-hydroxyethyl; R<sup>3</sup> is



R<sup>9</sup> is 2-(1R-hydroxyethyl-pyrimidin-4-yl), a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug.

82. A compound of claim 62, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, wherein R<sup>9</sup> is [1,3,5]-triazin-2-yl optionally substituted with up to two hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, morpholinyl or phenyl.

83. The compound of claim 82, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4-morpholin-4-yl-[1,3,5]-triazin-2-yl.

84. The compound of claim 82, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4-methoxy-6-methyl-[1,3,5]-triazin-2-yl.

85. The compound of claim 82, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4,6-dimethoxy-[1,3,5]-triazin-2-yl.

86. A compound of claim 82, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4-phenyl-[1,3,5]-triazin-2-yl.

87. A compound of claim 82, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4-cyclopropyl-[1,3,5]-triazin-2-yl.

88. A compound of claim 82, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4,6-dimethyl-[1,3,5]-triazin-2-yl.

89. A compound of claim 82, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4-hydroxymethyl-6-phenyl-[1,3,5]-triazin-2-yl.

90. A compound of claim 82, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4-methoxy-6-methoxymethyl-[1,3,5]-triazin-2-yl.

91. A compound of claim 82, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4-methyl-[1,3,5]-triazin-2-yl.


92. A compound of claim 82, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 4-methoxymethyl-6-phenyl-[1,3,5]triazin-2-yl.

93. The compound of claim 83, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, selected from 1R-{4-[2R,6S-dimethyl-4-(4-morpholin-4-yl-[1,3,5]triazin-2-yl)-piperazin-1-yl]-pyrimidin-2-yl}-ethanol; 1R-{4-[4-(4-methoxy-6-methyl-[1,3,5]triazin-2-yl)-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol; 1R-{4-[4-(4,6-dimethoxy-[1,3,5]triazin-2-yl)-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol; 1R-{4-[4-(4-methoxymethyl-6-phenyl-[1,3,5]triazin-2-yl)-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol; and 1R-{4-[2R,6S-dimethyl-4-(4-phenyl-[1,3,5]triazin-2-yl)-piperazin-1-yl]-pyrimidin-2-yl}-ethanol.

94. A compound of claim 13, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein:

R<sup>1</sup> is (R)-1-hydroxy-ethyl;

R<sup>3</sup> is



; and

R<sup>9</sup> is pyrimidyl, quinoxalyl or oxazolopyridyl optionally substituted with up to two (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy or hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl.

95. The compound of claim 94, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein:  $R^9$  is 4-hydroxymethyl-6-methyl-pyrimid-2-yl.

96. The compound of claim 94, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein:  $R^9$  is 2-hydroxymethyl-pyrimid-4-yl.

97. The compound of claim 94, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein:  $R^9$  is 2-hydroxymethyl-6-methyl-pyrimid-4-yl.

98. The compound of claim 94, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein:  $R^9$  is 2-(oxazolo[5,4-b]pyridyl).

99. The compound of claim 94, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein:  $R^9$  is 2-(oxazolo[4,5-b]pyridyl).


100. The compound of claim 94, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein:  $R^9$  is 2-quinoxalyl.

101. A compound of claim 94, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, selected from 1R-{4-[4-(4-hydroxymethyl-6-methyl-pyrimidin-2-yl)-3S-methyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol; 1R-{4-[4-(2-hydroxymethyl-pyrimidin-4-yl)-3S-methyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol; 1R-{4-[4-(2-hydroxymethyl-6-methyl-pyrimidin-4-yl)-3S-methyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol; 1R-[4-(3S-methyl-4-oxazolo[5,4-b]pyridin-2-yl)-piperazin-1-yl]-pyrimidin-2-yl]-ethanol; 1R-[4-(3S-methyl-4-oxazolo[4,5-b]pyridin-2-yl)-piperazin-1-yl]-pyrimidin-2-yl]-ethanol; and 1R-[4-(3S-methyl-4-quinoxalin-2-yl)-piperazin-1-yl]-pyrimidin-2-yl]-ethanol.

102. A compound of claim 13, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein:

$R^1$  is (R)-1-hydroxy-ethyl;

$R^3$  is



; and

R<sup>9</sup> is pyrimidyl optionally substituted with up to two (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl.


103. The compound of claim 102, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 2-(1R-hydroxyethyl)-pyrimid-4-yl.

104. The compound of claim 102 which is 1R-(4-{4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-2R-methyl-piperazin-1-yl}-pyrimidin-2-yl)-ethanol.

105. A compound of claim 13, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein:

R<sup>1</sup> is (R)-1-hydroxy-ethyl;

R<sup>3</sup> is



R<sup>9</sup> is pyrimidyl optionally substituted with up to two (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl.


106. The compound of claim 105, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 2-(1R-hydroxyethyl)-pyrimid-4-yl.

107. The compound of claim 105, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, which is (4-{4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-2R,5S-dimethyl-piperazin-1-yl}-pyrimidin-2-yl)-ethanol.

108. A compound of claim 13, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein:

R<sup>1</sup> is (S)-1-hydroxy-ethyl;

R<sup>3</sup> is



R<sup>9</sup> is pyrimidyl optionally substituted with up to two (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy or hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl.


109. The compound of claim 108, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 2-(1R-hydroxy-ethyl)-pyrimid-4-yl.

110. The compound of claim 108, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, which is 1S-(4-{4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl}-pyrimidin-2-yl)-ethanol.

111. A compound of claim 13, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein:

R<sup>1</sup> is acetyl;

R<sup>3</sup> is



R<sup>9</sup> is pyrimidyl optionally substituted with up to two (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, acetyl or hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl.

112. The compound of claim 111, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 2-acetyl-pyrimid-4-yl.

113. The compound of claim 111, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein: R<sup>9</sup> is 2-(1R-hydroxyethyl)-pyrimid-4-yl.

114. The compound of claim 111 which is 1-{4-[4-(2-acetyl-pyrimidin-4-yl)-2R\*,6S\*-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanone or 1-(-4-{4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl}-pyrimidin-2-yl)-ethanone.

115. A pharmaceutical composition comprising a compound of claim 1, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug and a pharmaceutically acceptable carrier or diluent.

116. A method of inhibiting sorbitol dehydrogenase in a mammal in need of such inhibition comprising administering to said mammal a sorbitol dehydrogenase inhibiting amount of a compound of claim 1, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug.



117. A method of treating diabetes in a mammal suffering from diabetes comprising administering to said mammal an effective amount of a compound of claim 1, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug.

5 118. A method of treating or preventing diabetic complications in a mammal comprising administering to said mammal an effective amount of a compound of claim 1, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug.

10 119. A method of claim 118 wherein said mammal is suffering from diabetes.

120. A method of claim 118 wherein said diabetic complication is diabetic neuropathy.

121. A method of claim 118 wherein said diabetic complication is diabetic nephropathy.

15 122. A method of claim 118 wherein said diabetic complication is diabetic retinopathy.

123. A method of claim 118 wherein said diabetic complication is foot ulcers.

20 124. A method of claim 118 wherein said diabetic complication is a cardiovascular condition.

125. A pharmaceutical composition comprising a compound of claim 1, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug and an aldose reductase inhibitor, a prodrug thereof or a pharmaceutically acceptable salt of said aldose reductase inhibitor or said prodrug.


25 126. A composition of claim 125 additionally comprising a pharmaceutically acceptable carrier or diluent.

127. A composition of claim 126 wherein said aldose reductase inhibitor is selected from the group consisting of ponalrestat, tolrestat, epalrestat, zenarestat, 2R,4R-6,7-dichloro-4-hydroxy-2-methylchroman-4-acetic acid, 2R,4R-6,7-dichloro-6-fluoro-4-hydroxy-2-methylchroman-4-acetic acid, 3,4-dihydro-2,8-diisopropyl-3-oxo-2H-1,4-benzoxazine-4-acetic acid, 3,4-dihydro-3-oxo-4-[(4,5,7-trifluoro-2-benzothiazolyl)methyl]-2H-1,4-benzothiazine-2-acetic acid, N-[3,5-dimethyl-4-[(nitromethyl)sulfonyl]phenyl]-2-methyl-benzeneacetamide, (S)-6-fluorospiro[chroman-4,4'-imidazolidine]-2,5'-dione, d-2-methyl-6-fluoro-spiro(chroman-4',4'-imidazolidine)-2',5'-dione, 2-fluoro-spiro(9H-fluorene-

30  
35

9,4'-imidazolidine)-2',5'-dione, 2,7-difluoro-spiro(9H-fluorene-9,4'-imidazolidine)-2',5'-dione, 2,7-difluoro-5-methoxy-spiro(9H-fluorene-9,4'-imidazolidine)-2',5'-dione, 7-fluoro-spiro(5H-indenol[1,2-b]pyridine-5,3'-pyrrolidine)-2,5'-dione, -cis-6'-chloro-2',3'-dihydro-2'-methyl-spiro-(imidazolidine-4,4'-4'-H-pyrano(2,3-b)pyridine)-2,5-dione, spiro[imidazolidine-4,5'-(6H)-quinoline]-2,5-dione-3'-chloro-7,'8'-dihydro-7'-methyl-(5'-cis), (2S,4S)-6-fluoro-2',5'-dioxospiro(chroman-4,4'-imidazolidine)-2-carboxamide, and 2-[(4-bromo-2-fluorophenyl)methyl]-6-fluorospiro[isoquinoline-4-(1H),3'-pyrrolidine]-1,2',3,5'(2H)-tetrone.

128. A composition of claim 126 wherein said aldose reductase inhibitor is selected from the group consisting of compounds of the formula ARI,



ARI

or a pharmaceutically acceptable salt thereof, wherein

Z in the compound of formula ARI is O or S;

R<sup>1</sup> in the compound of formula ARI is hydroxy or a group capable of being removed *in vivo* to produce a compound of formula ARI wherein R<sup>1</sup> is OH; and

X and Y in the compound of formula ARI are the same or different and are selected from hydrogen, trifluoromethyl, fluoro, and chloro.

129. A composition of claim 128 wherein said aldose reductase inhibitor is selected from the group consisting of 3,4-dihydro-3-(5-fluorobenzothiazol-2-ylmethyl)-4-oxophthalazin-1-yl-acetic acid, 3-(5,7-difluorobenzothiazol-2-ylmethyl)-3,4-dihydro-4-oxophthalazin-1-ylacetic acid, 3-(5-chlorobenzothiazol-2-ylmethyl)-3,4-dihydro-4-oxophthalazin-1-ylacetic acid, 3-(5,7-dichlorobenzothiazol-2-ylmethyl)-3,4-dihydro-4-oxophthalazin-1-ylacetic acid, 3-(5-trifluoromethylbenzoxazol-2-ylmethyl)phthalazin-1-ylacetic acid, 3,4-dihydro-3-(5-fluorobenzoxazol-2-ylmethyl)-4-oxophthalazin-1-yl-acetic acid, 3-(5,7-difluorobenzoxazol-2-ylmethyl)-3,4-dihydro-4-oxophthalazin-1-ylacetic acid, 3-(5-chlorobenzoxazol-2-ylmethyl)-3,4-dihydro-4-oxophthalazin-1-ylacetic acid, 3-(5,7-dichlorobenzoxazol-2-ylmethyl)-3,4-dihydro-4-oxophthalazin-1-ylacetic acid, and zopolrestat.

130. A method of treating diabetes in a mammal suffering from diabetes comprising administering to said mammal an effective amount of a compound of claim 1, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug and an aldose reductase inhibitor, a prodrug of said aldose reductase inhibitor or a pharmaceutically acceptable salt of said aldose reductase inhibitor or said prodrug.

131. A method of treating or preventing diabetic complications in a mammal comprising administering to said mammal an effective amount of a compound of claim 1, a prodrug thereof or a pharmaceutically acceptable of said compound or said prodrug and an aldose reductase inhibitor, a prodrug of said aldose reductase inhibitor or a pharmaceutically acceptable salt of said aldose reductase inhibitor or said prodrug thereof.

132. A method of claim 131 wherein said mammal is suffering from diabetes.

133. A method of claim 131 wherein said diabetic complication is diabetic neuropathy.

134. A method of claim 131 wherein said diabetic complication is diabetic nephropathy.

135. A method of claim 131 wherein said diabetic complication is diabetic retinopathy.


136. A method of claim 131 wherein said diabetic complication is foot ulcers.

137. A method of claim 131 wherein said diabetic complication is a cardiovascular condition.

138. A pharmaceutical composition comprising a compound of claim 1, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug and a sodium hydrogen ion exchange (NHE-1) inhibitor, a prodrug of said NHE-1 inhibitor or a pharmaceutically acceptable salt of said NHE-1 inhibitor or said prodrug thereof.

139. A composition of claim 138 additionally comprising a pharmaceutically acceptable carrier or diluent.

140. A composition of claim 138 wherein said NHE-1 inhibitor is a compound of the formula NHE,



NHE

a prodrug thereof or a pharmaceutically acceptable salt of said compound or of said prodrug, wherein

- 5           Z in the compound of formula NHE is carbon connected and is a five-membered, diaza, diunsaturated ring having two contiguous nitrogens, said ring optionally mono-, di-, or tri-substituted with up to three substituents independently selected from R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> ;

or

- 10           Z in the compound of formula NHE carbon connected and is a five-membered, triaza, diunsaturated ring, said ring optionally mono- or di-substituted with up to two substituents independently selected from R<sup>4</sup> and R<sup>5</sup>;

- 15           wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup> and R<sup>5</sup> in the compound of formula NHE are each independently hydrogen, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkylthio, (C<sub>3</sub>-C<sub>4</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, (C<sub>1</sub>-C<sub>4</sub>)alkoxy(C<sub>1</sub>-C<sub>4</sub>)alkyl, mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylcarbamoyl, M or M(C<sub>1</sub>-C<sub>4</sub>)alkyl, any of said previous (C<sub>1</sub>-C<sub>4</sub>)alkyl moieties optionally having from one to nine fluorines; said (C<sub>1</sub>-C<sub>4</sub>)alkyl or (C<sub>3</sub>-C<sub>4</sub>)cycloalkyl optionally
- 20           mono- or di-substituted independently with hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, (C<sub>1</sub>-C<sub>4</sub>)alkylthio, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfinyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl, mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylcarbamoyl or mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylaminosulfonyl; and said (C<sub>3</sub>-C<sub>4</sub>)cycloalkyl optionally having from one to seven fluorines;

- wherein M in the compound of formula NHE is a partially saturated,
- 25           fully saturated or fully unsaturated five to eight membered ring optionally having one to three heteroatoms selected independently from oxygen, sulfur and nitrogen, or, a bicyclic ring consisting of two fused partially saturated, fully saturated or fully unsaturated three to six membered rings, taken independently, optionally having one to four heteroatoms selected
- 30           independently from nitrogen, sulfur and oxygen;

              said M in the compound of formula NHE is optionally substituted, on one ring if the moiety is monocyclic, or one or both rings if the moiety is bicyclic, on carbon or nitrogen with up to three substituents independently

selected from R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup>, wherein one of R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> is optionally a partially saturated, fully saturated, or fully unsaturated three to seven membered ring optionally having one to three heteroatoms selected independently from oxygen, sulfur and nitrogen optionally substituted with

5 (C<sub>1</sub>-C<sub>4</sub>)alkyl and additionally R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are optionally hydroxy, nitro, halo, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl, formyl, (C<sub>1</sub>-C<sub>4</sub>)alkanoyl, (C<sub>1</sub>-C<sub>4</sub>)alkanoyloxy, (C<sub>1</sub>-C<sub>4</sub>)alkanoylamino, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonylamino, sulfonamido, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonamido, amino, mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylamino, carbamoyl, mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylcarbamoyl, cyano,  
10 thiol, (C<sub>1</sub>-C<sub>4</sub>)alkylthio, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfinyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonyl, mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylaminosulfonyl, (C<sub>2</sub>-C<sub>4</sub>)alkenyl, (C<sub>2</sub>-C<sub>4</sub>)alkynyl or (C<sub>5</sub>-C<sub>7</sub>)cycloalkenyl,

wherein said (C<sub>1</sub>-C<sub>4</sub>)alkoxy, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>7</sub>)alkanoyl, (C<sub>1</sub>-C<sub>4</sub>)alkylthio, mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylamino or (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> substituents  
15 are optionally mono- substituted independently with hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>1</sub>-C<sub>4</sub>)alkanoyl, (C<sub>1</sub>-C<sub>4</sub>)alkanoylamino, (C<sub>1</sub>-C<sub>4</sub>)alkanoyloxy, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonylamino, sulfonamido, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonamido, amino, mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylamino, carbamoyl, mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylcarbamoyl, cyano, thiol, nitro, (C<sub>1</sub>-C<sub>4</sub>)alkylthio, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfinyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonyl or mono-  
20 N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylaminosulfonyl or optionally substituted with one to nine fluorines.

141. A composition of claim 140 wherein said NHE-1 inhibitor is selected from the group consisting of [1-(8-bromoquinolin-5-yl)-5-cyclopropyl-1H-pyrazole-4-carbonyl]guanidine;

25 [1-(6-chloroquinolin-5-yl)-5-cyclopropyl-1H-pyrazole-4-carbonyl]guanidine; [1-(indazol-7-yl)-5-cyclopropyl-1H-pyrazole-4-carbonyl]guanidine;  
[1-(benzimidazol-5-yl)-5-cyclopropyl-1H-pyrazole-4-carbonyl]guanidine;  
[1-(1-isoquinolyl)-5-cyclopropyl-1H-pyrazole-4-carbonyl]guanidine;  
[5-cyclopropyl-1-(4-quinolinyl)-1H-pyrazole-4-carbonyl]guanidine;  
30 [5-cyclopropyl-1-(quinolin-5-yl)-1H-pyrazole-4-carbonyl]guanidine;  
[5-cyclopropyl-1-(quinolin-8-yl)-1H-pyrazole-4-carbonyl]guanidine;  
[1-(indazol-6-yl)-5-ethyl-1H-pyrazole-4-carbonyl]guanidine;  
[1-(indazol-5-yl)-5-ethyl-1H-pyrazole-4-carbonyl]guanidine;  
[1-(benzimidazol-5-yl)-5-ethyl-1H-pyrazole-4-carbonyl]guanidine;  
35 [1-(1-methylbenzimidazol-6-yl)-5-ethyl-1H-pyrazole-4-carbonyl]guanidine;

- 1-(5-quinoliny)-5-*n*-propyl-1*H*-pyrazole-4-carbonyl]guanidine;  
[1-(5-quinoliny)-5-isopropyl-1*H*-pyrazole-4-carbonyl]guanidine;  
[5-ethyl-1-(6-quinoliny)-1*H*-pyrazole-4-carbonyl]guanidine;  
[1-(2-methylbenzimidazol-5-yl)-5-ethyl-1*H*-pyrazole-4-carbonyl]guanidine; [1-  
5 (1,4-benzodioxan-6-yl)-5-ethyl-1*H*-pyrazole-4-carbonyl]guanidine;  
[1-(benzotriazol-5-yl)-5-ethyl-1*H*-pyrazole-4-carbonyl]guanidine;  
[1-(3-chloroindazol-5-yl)-5-ethyl-1*H*-pyrazole-4-carbonyl]guanidine;  
[1-(5-quinoliny)-5-butyl-1*H*-pyrazole-4-carbonyl]guanidine;  
[5-propyl-1-(6-quinoliny)-1*H*-pyrazole-4-carbonyl]guanidine;  
10 [5-isopropyl-1-(6-quinoliny)-1*H*-pyrazole-4-carbonyl]guanidine;  
[1-(2-chloro-4-methylsulfonylphenyl)-5-cyclopropyl-1*H*-pyrazole-4-  
carbonyl]guanidine;  
[1-(2-chlorophenyl)-5-cyclopropyl-1*H*-pyrazole-4-carbonyl]guanidine;  
[1-(2-trifluoromethyl-4-fluorophenyl)-5-cyclopropyl-1*H*-pyrazole-4-  
15 carbonyl]guanidine;  
[1-(2-bromophenyl)-5-cyclopropyl-1*H*-pyrazole-4-carbonyl]guanidine;  
[1-(2-fluorophenyl)-5-cyclopropyl-1*H*-pyrazole-4-carbonyl]guanidine;  
[1-(2-chloro-5-methoxyphenyl)-5-cyclopropyl-1*H*-pyrazole-4-  
carbonyl]guanidine;  
20 [1-(2-chloro-4-methylaminosulfonylphenyl)-5-cyclopropyl-1*H*-pyrazole-4-  
carbonyl]guanidine;  
[1-(2,5-dichlorophenyl)-5-cyclopropyl-1*H*-pyrazole-4-carbonyl]guanidine;  
[1-(2,3-dichlorophenyl)-5-cyclopropyl-1*H*-pyrazole-4-carbonyl]guanidine;  
[1-(2-chloro-5-aminocarbonylphenyl)-5-cyclopropyl-1*H*-pyrazole-4-  
25 carbonyl]guanidine;  
[1-(2-chloro-5-aminosulfonylphenyl)-5-cyclopropyl-1*H*-pyrazole-4-  
carbonyl]guanidine;  
[1-(2-fluoro-6-trifluoromethylphenyl)-5-cyclopropyl-1*H*-pyrazole-4-  
carbonyl]guanidine;  
30 [1-(2-chloro-5-methylsulfonylphenyl)-5-cyclopropyl-1*H*-pyrazole-4-  
carbonyl]guanidine;  
[1-(2-chloro-5-dimethylaminosulfonylphenyl)-5-cyclopropyl-1*H*-pyrazole-4-  
carbonyl]guanidine;  
[1-(2-trifluoromethyl-4-chlorophenyl)-5-cyclopropyl-1*H*-pyrazole-4-  
35 carbonyl]guanidine;

[1-(2-chlorophenyl)-5-methyl-1H-pyrazole-4-carbonyl]guanidine;  
[5-methyl-1-(2-trifluoromethylphenyl)-1H-pyrazole-4-carbonyl]guanidine;  
[5-ethyl-1-phenyl-1H-pyrazole-4-carbonyl]guanidine;  
[5-cyclopropyl-1-(2-trifluoromethylphenyl)-1H-pyrazole-4-carbonyl]guanidine;  
5 [5-cyclopropyl-1-phenyl-1H-pyrazole-4-carbonyl]guanidine;  
[5-cyclopropyl-1-(2,6-dichlorophenyl)-1H-pyrazole-4-carbonyl]guanidine; or a  
pharmaceutically acceptable salt thereof.

142. A method of treating ischemia in a mammal suffering from ischemia  
comprising administering to said mammal an effective amount of a compound of  
10 claim 1, a prodrug thereof or a pharmaceutically acceptable salt of said compound or  
said prodrug and a sodium hydrogen ion exchange (NHE-1) inhibitor, a prodrug of  
said NHE-1 inhibitor or a pharmaceutically acceptable salt of said NHE-1 inhibitor or  
said prodrug.

143. A method of claim 142 wherein said ischemia is perioperative  
15 myocardial ischemia.

144. A method of treating or preventing diabetic complications in a  
mammal comprising administering to said mammal an effective amount of a  
compound of claim 1, a prodrug thereof or a pharmaceutically acceptable salt of said  
compound or said prodrug and a sodium hydrogen ion exchange (NHE-1) inhibitor, a  
20 prodrug of said NHE-1 inhibitor or a pharmaceutically acceptable salt of said NHE-1  
inhibitor or said prodrug.

145. A method of claim 144 wherein said mammal is suffering from  
diabetes.

146. A method of claim 144 wherein said diabetic complication is diabetic  
25 neuropathy.

147. A method of claim 144 wherein said diabetic complication is diabetic  
nephropathy.

148. A method of claim 144 wherein said diabetic complication is diabetic  
retinopathy.

30 149. A method of claim 144 wherein said diabetic complication is foot  
ulcers.

150. A method of claim 144 wherein said diabetic complication is a  
cardiovascular condition.

151. A method of treating diabetes in a mammal suffering from diabetes  
35 comprising administering to said mammal an effective amount of a compound of

claim 1, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, and a sodium hydrogen ion exchange (NHE-1) inhibitor, a prodrug of said NHE-1 inhibitor or a pharmaceutically acceptable salt of said NHE-1 inhibitor or said prodrug.

5           152. A kit comprising:

          a. a compound of claim 1, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug in a first unit dosage form;

          b. an aldose reductase inhibitor, a prodrug thereof or a pharmaceutically acceptable salt of said prodrug or said aldose reductase inhibitor in a second unit dosage form; and

          c. a container.

          153. A kit comprising:

15           a. a compound of claim 1, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug in a first unit dosage form;

          b. a sodium hydrogen ion exchange (NHE-1) inhibitor, a prodrug thereof or a pharmaceutically acceptable salt of said prodrug or said NHE-1 inhibitor in a second unit dosage form; and

20           c. a container.

          154. A method of inhibiting sorbitol dehydrogenase in a mammal in need thereof comprising administering to said mammal a pharmaceutical composition of claim 115.

25           155. A method of treating ischemia in a mammal suffering from ischemia comprising administering to said mammal a pharmaceutical composition of claim 138.

          156. A method of claim 154 wherein said ischemia is perioperative myocardial ischemia.

30           157. A method of treating or preventing diabetic complications in a mammal comprising administering to said mammal a pharmaceutical composition of claim 115.

          158. A method of claim 157 wherein said mammal is suffering from diabetes.



159. A method of treating or preventing diabetic complications in a mammal comprising administering to said mammal a pharmaceutical composition of claim 125.

160. A method of claim 159 wherein said mammal is suffering from diabetes.

161. A method of treating or preventing diabetic complications in a mammal comprising administering to said mammal a pharmaceutical composition of claim 138.


162. A method of claim 161 wherein said mammal is suffering from diabetes.

163. A compound of claim 1, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, wherein:


$R^1$  is  $C(OH)R^4R^5$ , where  $R^4$  and  $R^5$  are each independently hydrogen or methyl;

$R^2$  is hydrogen;

$R^3$  is



or



wherein said piperazinyl  $R^3$  is substituted by  $R^6$ ,  $R^7$  or  $R^8$ ;

$G$ ,  $G^1$  and  $G^2$  are taken separately and are each hydrogen and  $R^6$  is hydrogen or  $(C_1-C_4)$ alkyl;  $R^7$  and  $R^8$  are each independently hydrogen or  $(C_1-C_4)$ alkyl; or

$G$  and  $G^1$  are taken together and are  $(C_1-C_3)$ alkylene and  $R^6$ ,  $R^7$ ,  $R^8$  and  $G^2$  are hydrogen; or

$G^1$  and  $G^2$  are taken together and are  $(C_1-C_3)$ alkylene and  $R^6$ ,  $R^7$ ,  $R^8$  and  $G$  are hydrogen;

$q$  is 0 or 1;

$X$  is a covalent bond, oxycarbonyl, vinylenylcarbonyl, oxy $(C_1-C_4)$ alkylenylcarbonyl, thio $(C_1-C_4)$ alkylenylcarbonyl or vinylenylsulfonyl; said vinylenylcarbonyl and said vinylenylsulfonyl in the definition of  $X$  are optionally substituted on one or two vinylenyl carbons with  $(C_1-C_4)$ alkyl, benzyl or Ar; said oxy $(C_1-C_4)$ alkylenylcarbonyl

and said thio(C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl in the definition of X are optionally substituted with up to two (C<sub>1</sub>-C<sub>4</sub>)alkyl, benzyl or Ar;

R<sup>9</sup> is (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, Ar<sup>1</sup>-(C<sub>0</sub>-C<sub>4</sub>)alkylenyl or (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted with up to five fluoro;

- 5 Ar<sup>1</sup> is phenyl, naphthyl, pyridyl, pyrimidyl, pyrazinyl, pyridazinyl, triazinyl, quinolyl, isoquinolyl, quinazolyl, quinoxalyl, phthalazinyl, cinnolyl, naphthyridinyl, pteridinyl, pyrazinopyrazinyl, pyrazinopyridazinyl, pyrimidopyridazinyl, pyrimidopyrimidyl, pyridopyrimidyl, pyridopyrazinyl, pyridopyridazinyl, pyrrolyl, furanyl, thienyl, imidazolyl, oxazolyl, thiazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzoxazolyl, benzothiazolyl, indazolyl, benzisoxazolyl, benzisothiazolyl, pyrrolopyridyl, furopyridyl, thienopyridyl, imidazolopyridyl, oxazolopyridyl, thiazolopyridyl, pyrazolopyridyl, isoxazolopyridyl, isothiazolopyridyl, pyrrolopyrimidyl, furopyrimidyl, thienopyrimidyl, imidazolopyrimidyl, oxazolopyrimidyl, thiazolopyrimidyl, pyrazolopyrimidyl, isoxazolopyrimidyl, isothiazolopyrimidyl, pyrrolopyrazinyl, furopyrazinyl, thienopyrazinyl, imidazolopyrazinyl, oxazolopyrazinyl, thiazolopyrazinyl, pyrazolopyrazinyl, isoxazolopyrazinyl, isothiazolopyrazinyl, pyrrolopyridazinyl, furopyridazinyl, thienopyridazinyl, imidazolopyridazinyl, oxazolopyridazinyl, thiazolopyridazinyl, pyrazolopyridazinyl, isoxazolopyridazinyl or isothiazolopyridazinyl;
- 10  
15  
20 and

said Ar<sup>1</sup> is optionally substituted as set forth above;

k is 0, 1, 2, 3 or 4;

Y<sup>1</sup> is a covalent bond, carbonyl, sulfonyl or oxycarbonyl;

- 25 R<sup>43</sup> is (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, Ar<sup>5</sup>-(C<sub>0</sub>-C<sub>4</sub>)alkylenyl, NR<sup>47</sup>R<sup>48</sup> or (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted with one to five fluoro; provided that when Y<sup>1</sup> is a covalent bond or oxycarbonyl, then R<sup>43</sup> is not NR<sup>47</sup>R<sup>48</sup>;

R<sup>47</sup> and R<sup>48</sup> are taken separately and are each independently selected from hydrogen, Ar<sup>5</sup>, (C<sub>1</sub>-C<sub>6</sub>)alkyl and Ar<sup>5</sup>-(C<sub>0</sub>-C<sub>4</sub>)alkylenyl; or

- 30 R<sup>47</sup> and R<sup>48</sup> are taken together with the nitrogen atom to which they are attached to form azetidiny, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, azepinyl, azabicyclo[3.2.2]nonanyl, azabicyclo[2.2.1]heptyl, 1,2,3,4-tetrahydroisoquinolyl, 6,7-dihydro-5H-dibenzo[c,e]azepinyl or 5,6,7,8-tetrahydropyrido[4,3-d]pyrimidyl; said azetidiny in the definition of R<sup>47</sup> and R<sup>48</sup> are optionally substituted with one hydroxy, amino, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to
- 35

fluoro; said pyrrolidinyl, piperidinyl and azepinyl in the definition of R<sup>47</sup> and R<sup>48</sup> are optionally substituted with up to two hydroxy, amino, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said morpholinyl in the  
 5 definition of R<sup>47</sup> and R<sup>48</sup> is optionally substituted with up to two substituents independently selected from hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said piperazinyl, 1,2,3,4-tetrahydroisoquinolyl and 5,6,7,8-tetrahydro[4,3-d]pyrimidyl in the definition of R<sup>47</sup> and R<sup>48</sup> are optionally  
 10 substituted with up to three hydroxy, amino, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; and said 6,7-dihydro-5H-dibenzo[c,e]azepinyl in the definition of R<sup>47</sup> and R<sup>48</sup> are optionally substituted with up to four hydroxy, amino, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally  
 15 substituted with up to five fluoro;

Ar<sup>5</sup> is independently defined as set forth for Ar and Ar<sup>1</sup> above;

Ar<sup>5</sup> is optionally independently substituted as set forth for Ar and Ar<sup>1</sup> above.

164. A compound selected from

20 1R-(4-{1'-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-[4,4']bipiperidinyl-1-yl}-pyrimidin-2-yl)-ethanol;  
 furo[3,2-c]pyridin-2-yl-{4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-3R,5S-dimethyl-piperazin-1-yl}-methanone;  
 (4-chloro-furo[3,2-c]pyridin-2-yl)-{4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-3R,5S-  
 25 dimethyl-piperazin-1-yl}-methanone;  
 {4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-3R,5S-dimethyl-piperazin-1-yl}-(4-pyrrolidin-1-yl-furo[3,2-c]pyridin-2-yl)-methanone;  
 {4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-3R,5S-dimethyl-piperazin-1-yl}-(4-morpholin-4-yl-furo[3,2-c]pyridin-2-yl)-methanone;  
 30 {4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-3R,5S-dimethyl-piperazin-1-yl}-imidazo[1,2-a]pyridin-2-yl-methanone;  
 furo[3,2-c]pyridin-2-yl-{4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl}-methanone;  
 4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazine-1-carboxylic acid  
 35 pyridin-3-yl ester;

- 4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazine-1-carboxylic acid  
2-methyl-pyridin-3-yl ester;  
4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazine-1- carboxylic acid  
5-chloro-pyridin-3-yl ester;
- 5 4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazine-1-carboxylic acid  
6-methyl-pyridin-3-yl ester;  
(E)-1-{4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl}-3-  
thiophen-2-yl-propenone;  
1R-{4-[4-(4,6-dimethyl-pyrimidin-2-yl)-3R,5S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-  
10 ethanol;  
1R-{4-[4-(4-methoxymethyl-6-methyl-pyrimidin-2-yl)-3R,5S-dimethyl-piperazin-1-yl]-  
pyrimidin-2-yl}-ethanol;  
1R-{4-[4-(4-hydroxymethyl-6-methyl-pyrimidin-2-yl)-3R,5S-dimethyl-piperazin-1-yl]-  
pyrimidin-2-yl}-ethanol;
- 15 1R-(4-{3R,5S-dimethyl-4-[2-(4-methyl-piperazin-1-yl)-pyrimidin-4-yl]-piperazin-1-yl}-  
pyrimidin-2-yl)-ethanol;  
1R-(4-{4-[2-(4-ethyl-piperazin-1-yl)-pyrimidin-4-yl]-3R,5S-dimethyl-piperazin-1-yl}-  
pyrimidin-2-yl)-ethanol;  
1R-(4-{3R,5S-dimethyl-4-[2-(4-methyl-imidazol-1-yl)-pyrimidin-4-yl]-piperazin-1-yl}-  
20 pyrimidin-2-yl)-ethanol;  
1R-(4-{3R,5S-dimethyl-4-[2-(2-methyl-imidazol-1-yl)-pyrimidin-4-yl]-piperazin-1-yl}-  
pyrimidin-2-yl)-ethanol;  
1R-(4-{4-[2-(2,4-dimethyl-imidazol-1-yl)-pyrimidin-4-yl]-3R,5S-dimethyl-piperazin-1-  
yl}-pyrimidin-2-yl)-ethanol;
- 25 1R-(4-{4-[2-(4-isopropyl-piperazin-1-yl)-pyrimidin-4-yl]-3R,5S-dimethyl-piperazin-1-  
yl}-pyrimidin-2-yl)-ethanol;  
1R-(4-{3R,5S-dimethyl-4-[4-methyl-6-(4-methyl-piperazin-1-yl)-[1,3,5]triazin-2-yl]-  
piperazin-1-yl}-pyrimidin-2-yl)-ethanol;  
1R-{4-[4-(4-methoxy-6-methyl-[1,3,5]triazin-2-yl)-3R,5S-dimethyl-piperazin-1-yl]-  
30 pyrimidin-2-yl}-ethanol;  
1R-{4-[4-(4,6-dimethoxy-[1,3,5]triazin-2-yl)-3R,5S-dimethyl-piperazin-1-yl]-pyrimidin-  
2-yl}-ethanol;  
1R-{4-[4-(4-ethoxy-6-methyl-[1,3,5]triazin-2-yl)-3R,5S-dimethyl-piperazin-1-yl]-  
pyrimidin-2-yl}-ethanol;

1R-{4-[4-(4-isopropoxy-6-methyl-[1,3,5]triazin-2-yl)-3R,5S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;

1R-{4-[3R,5S-dimethyl-4-(4-phenyl-[1,3,5]triazin-2-yl)-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;

5 1R-{4-[4-(4-hydroxymethyl-6-methoxy-[1,3,5]triazin-2-yl)-3R,5S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;

1R-{4-[4-(4-isopropoxy-6-methoxy-[1,3,5]triazin-2-yl)-3R,5S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;

10 1R-{4-[4-(4-isopropyl-[1,3,5]triazin-2-yl)-3R,5S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;

1R-{4-[4-(4-ethyl-6-methoxy-[1,3,5]triazin-2-yl)-3R,5S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;

1R-{4-[4-(4,6-dimethyl-pyrimidin-2-yl)-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;

15 1R-{4-[4-(4-hydroxymethyl-6-methyl-pyrimidin-2-yl)-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;

1R-{4-[2R,6S-dimethyl-4-(4-[1,2,4]triazol-1-yl-pyrimidin-2-yl)-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;

20 1R-{4-[4-(2,6-dimethyl-pyrimidin-4-yl)-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;

1R-{4-[4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;

1R-{4-[4-(2-hydroxymethyl-6-methyl-pyrimidin-4-yl)-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;

25 1R-{4-[4-[2-(1S-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;

1S-{4-[4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;

30 1-{4-[4-(2-acetyl-pyrimidin-4-yl)-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanone;

1RS-{4-[4-[2-(1RS-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;

(4-{4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-3R,5S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanone;

- 1R-{4-[2R,6S-dimethyl-4-(2-morpholin-4-yl-pyrimidin-4-yl)-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;
- 1R-(4-{2R,6S-dimethyl-4-[2-(4-methyl-piperazin-1-yl)-pyrimidin-4-yl]-piperazin-1-yl}-pyrimidin-2-yl)-ethanol;
- 5 1R-{4-[2R,6S-dimethyl-4-(2-[1,2,4]triazol-1-yl-pyrimidin-4-yl)-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;
- 1R-(4-{4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6R-dimethyl-piperazin-1-yl}-pyrimidin-2-yl)-ethanol;
- 1R-(4-{4-[2-(4-ethyl-piperazin-1-yl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl}-pyrimidin-2-yl)-ethanol;
- 10 1R-(4-{2R,6S-dimethyl-4-[2-(4-methyl-imidazol-1-yl)-pyrimidin-4-yl]-piperazin-1-yl}-pyrimidin-2-yl)-ethanol;
- 1R-(4-{4-[2-(2,4-dimethyl-imidazol-1-yl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl}-pyrimidin-2-yl)-ethanol;
- 15 1R-{4-[2R,6S-dimethyl-4-(4-morpholin-4-yl-[1,3,5]triazin-2-yl)-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;
- 1R-{4-[4-(4-methoxy-6-methyl-[1,3,5]triazin-2-yl)-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;
- 1R-{4-[4-(4,6-dimethoxy-[1,3,5]triazin-2-yl)-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;
- 20 1R-{4-[2R,6S-dimethyl-4-(4-phenyl-[1,3,5]triazin-2-yl)-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;
- 1R-{4-[4-(4-hydroxymethyl-6-methyl-pyrimidin-2-yl)-3S-methyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;
- 25 1R-{4-[4-(2-hydroxymethyl-pyrimidin-4-yl)-3S-methyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;
- 1R-{4-[4-(2-hydroxymethyl-6-methyl-pyrimidin-4-yl)-3S-methyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;
- 1R-{4-(3S-methyl-4-oxazolo[5,4-b]pyridin-2-yl)-piperazin-1-yl}-pyrimidin-2-yl}-ethanol;
- 30 1R-{4-(3S-methyl-4-oxazolo[4,5-b]pyridin-2-yl)-piperazin-1-yl}-pyrimidin-2-yl}-ethanol;
- 1R-{4-(3S-methyl-4-quinoxalin-2-yl)-piperazin-1-yl}-pyrimidin-2-yl}-ethanol;
- 1R-{4-[4-(4,6-dimethyl-[1,3,5]triazin-2-yl)-3R,5S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;
- 1R-{4-[3R,5S-dimethyl-4-(4-methyl-6-phenyl-[1,3,5]triazin-2-yl)-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;
- 35

1R-{4-[4-(4-cyclopropyl-[1,3,5]triazin-2-yl)-3R,5S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;

1R-{4-[4-(4-cyclopropyl-[1,3,5]triazin-2-yl)-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;

5 1R-{4-[4-(4,6-dimethyl-[1,3,5]triazin-2-yl)-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;

1R-{4-[4-(4-hydroxymethyl-6-phenyl-[1,3,5]triazin-2-yl)-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;

10 1R-{4-[4-(4-methoxy-6-methoxymethyl-[1,3,5]triazin-2-yl)-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;

1R-{4-[2R,6S-dimethyl-4-(4-methyl-[1,3,5]triazin-2-yl)-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;

1-{4-[4-(2-acetyl-pyrimidin-4-yl)-2R\*,6S\*-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanone;


15 1-(-4-{4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl)-ethanone;

1R-{4-[4-(4-methoxymethyl-6-phenyl-[1,3,5]-triazin-2-yl)-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethanol;

20 (4-{4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-2R,5S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl)-ethanol; and

1S-(4-{4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl)-ethanol.

165. A compound of the formula I<sup>A</sup>,




wherein:


R<sup>1</sup> is C-(OR<sup>80</sup>)R<sup>4</sup>R<sup>5</sup>, where R<sup>80</sup> is independently (C<sub>1</sub>-C<sub>4</sub>)alkyl, benzyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl or phenylcarbonyl, where said benzyl and said phenyl are optionally substituted with up to three (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, halo or nitro;

30 R<sup>4</sup> and R<sup>5</sup> are each independently hydrogen, methyl, ethyl or hydroxy-(C<sub>1</sub>-C<sub>3</sub>)alkyl; R<sup>2</sup> is hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl or (C<sub>1</sub>-C<sub>4</sub>)alkoxy;

R<sup>3</sup> is a radical of the formula







said radical of formula  $R^{3b}$  is substituted by  $R^{18}$ ,  $R^{19}$  and  $R^{20}$ ;

G,  $G^1$  and  $G^2$  are taken separately and are each hydrogen and  $R^6$  is hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl or phenyl optionally independently substituted with up to three hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl or (C<sub>1</sub>-C<sub>4</sub>)alkoxy, wherein said (C<sub>1</sub>-C<sub>4</sub>)alkyl in the definition of  $R^6$  and said (C<sub>1</sub>-C<sub>4</sub>)alkoxy in the definition of  $R^6$  are optionally and independently substituted with up to five fluoro;  $R^7$  and  $R^8$  are each independently hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl; or

G and  $G^1$  are taken together and are (C<sub>1</sub>-C<sub>3</sub>)alkylene and  $R^6$ ,  $R^7$ ,  $R^8$  and  $G^2$  are hydrogen; or

$G^1$  and  $G^2$  are taken together and are (C<sub>1</sub>-C<sub>3</sub>)alkylene and  $R^6$ ,  $R^7$ ,  $R^8$  and G are hydrogen;

q is 0 or 1;

X is a covalent bond, -(C=NR<sup>10</sup>)-, oxycarbonyl, vinylenylcarbonyl, oxy(C<sub>1</sub>-

C<sub>4</sub>)alkylenylcarbonyl, (C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl, (C<sub>3</sub>-C<sub>4</sub>)alkenylcarbonyl, thio(C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl, vinylenylsulfonyl, sulfinyl-(C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl, sulfonyl-(C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl or carbonyl(C<sub>0</sub>-C<sub>4</sub>)alkylenylcarbonyl; wherein said oxy(C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl, (C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl, (C<sub>3</sub>-C<sub>4</sub>)alkenylcarbonyl and thio(C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl in the definition of X are each optionally and independently substituted with up to two (C<sub>1</sub>-C<sub>4</sub>)alkyl, benzyl or Ar; said vinylenylsulfonyl and said vinylenylcarbonyl in the definition of X are optionally substituted independently on one or two vinylenyl carbons with (C<sub>1</sub>-C<sub>4</sub>)alkyl, benzyl or Ar; and said carbonyl(C<sub>0</sub>-C<sub>4</sub>)alkylenylcarbonyl in the definition of X is optionally substituted independently with up to three (C<sub>1</sub>-C<sub>4</sub>)alkyl, benzyl or Ar;

$R^{10}$  is hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl;

$R^9$  is (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, Ar<sup>1</sup>-(C<sub>0</sub>-C<sub>3</sub>)alkylenyl or (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted with up to five fluoro; provided that when q = 0 and X is a covalent bond, oxycarbonyl or (C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl, then  $R^9$  is not (C<sub>1</sub>-C<sub>6</sub>)alkyl;

Ar and Ar<sup>1</sup> are independently a fully saturated, partially saturated or fully unsaturated five- to eight-membered ring optionally having up to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of two fused independently partially saturated, fully saturated or fully unsaturated five- to seven-membered rings, taken independently, optionally having up to four heteroatoms selected independently from nitrogen, sulfur and oxygen, or a tricyclic ring consisting of three fused independently partially saturated, fully saturated or fully

unsaturated five to seven membered rings, taken independently, optionally having up to four heteroatoms selected independently from nitrogen, sulfur and oxygen, said partially saturated, fully saturated ring or fully unsaturated monocyclic ring, bicyclic ring or tricyclic ring optionally having one or two oxo groups substituted on carbon or one or two oxo groups substituted on sulfur;

Ar and Ar<sup>1</sup> are optionally independently substituted on carbon or nitrogen, on one ring if the moiety is monocyclic, on one or both rings if the moiety is bicyclic, or on one, two or three rings if the moiety is tricyclic, with up to a total of four substituents independently selected from R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup>; wherein R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> are

each taken separately and are each independently halo, formyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylenyloxycarbonyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, C(OH)R<sup>15</sup>R<sup>16</sup>, naphthyl, phenyl, imidazolyl, pyridyl, triazolyl, morpholinyl, (C<sub>0</sub>-C<sub>4</sub>)alkylsulfamoyl, N-(C<sub>0</sub>-C<sub>4</sub>)alkylcarbamoyl, N,N-di-(C<sub>1</sub>-C<sub>4</sub>)alkylcarbamoyl, N-phenylcarbamoyl, N-(C<sub>1</sub>-C<sub>4</sub>)alkyl-N-phenylcarbamoyl, N,N-diphenyl carbamoyl, (C<sub>1</sub>-C<sub>4</sub>)alkylcarbonylamido, (C<sub>3</sub>-C<sub>7</sub>)cycloalkylcarbonylamido, phenylcarbonylamido, piperidinyl, pyrrolidinyl, piperazinyl, cyano, benzimidazolyl, amino, anilino, pyrimidyl, oxazolyl, isoxazolyl, tetrazolyl, thienyl, thiazolyl, benzothiazolyl, pyrrolyl, pyrazolyl, tetrahydroquinolyl, tetrahydroisoquinolyl, benzoxazolyl, pyridazinyl, pyridyloxy, pyridylsulfanyl, furanyl, 8-(C<sub>1</sub>-C<sub>4</sub>)alkyl-3,8-diaza[3.2.1]bicyclooctyl, 3,5-dioxo-1,2,4-triazinyl, phenoxy, thiophenoxy, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfanyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said naphthyl, phenyl, pyridyl, piperidinyl, benzimidazolyl, pyrimidyl, thienyl, benzothiazolyl, pyrrolyl, tetrahydroquinolyl, tetrahydroisoquinolyl, benzoxazolyl, pyridazinyl, pyridyloxy, pyridylsulfanyl, furanyl, thiophenoxy, anilino and phenoxy in the definition of R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> are optionally substituted with up to three substituents independently selected from hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said imidazolyl, oxazolyl, isoxazolyl, thiazolyl and pyrazolyl in the definition of R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> are optionally substituted with up to two substituents independently selected from hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said morpholinyl in the definition of R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> is optionally substituted with up to two substituents independently selected from (C<sub>1</sub>-C<sub>4</sub>)alkyl; said pyrrolidinyl in the definition of R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> is

optionally substituted with up to two substituents independently selected from hydroxy, hydroxy-(C<sub>1</sub>-C<sub>3</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said piperazinyl in the definition of R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> is optionally substituted with up to three substituents independently selected from (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, hydroxy-(C<sub>1</sub>-C<sub>3</sub>)alkyl, phenyl, pyridyl, (C<sub>0</sub>-C<sub>4</sub>)alkylsulfamoyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said triazolyl in the definition of R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> is optionally substituted with hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said tetrazolyl in the definition of R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> is optionally substituted with hydroxy-(C<sub>2</sub>-C<sub>3</sub>)alkyl or (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro; and said phenyl and pyridyl which are optionally substituted on piperazine in the definition of R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> are optionally substituted with up to three hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; or R<sup>11</sup> and R<sup>12</sup> are taken together on adjacent carbon atoms and are -CH<sub>2</sub>OC(CH<sub>3</sub>)<sub>2</sub>OCH<sub>2</sub>- or -O-(CH<sub>2</sub>)<sub>p</sub>-O-, and R<sup>13</sup> and R<sup>14</sup> are taken separately and are each independently hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl; p is 1, 2 or 3; R<sup>15</sup> and R<sup>16</sup> are taken separately and are each independently hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro; or R<sup>15</sup> and R<sup>16</sup> are taken separately and R<sup>15</sup> is hydrogen and R<sup>16</sup> is (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, hydroxy-(C<sub>1</sub>-C<sub>3</sub>)alkyl, phenyl, pyridyl, pyrimidyl, thienyl, furanyl, thiazolyl, oxazolyl, imidazolyl, benzothiazolyl or benzoxazolyl; or R<sup>15</sup> and R<sup>16</sup> are taken together and are (C<sub>3</sub>-C<sub>6</sub>)alkylene; G<sup>3</sup>, G<sup>4</sup> and G<sup>5</sup> are taken separately and are each hydrogen; r is 0; R<sup>18</sup> is hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl or phenyl optionally independently substituted with up to three hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl or (C<sub>1</sub>-C<sub>4</sub>)alkoxy, wherein said (C<sub>1</sub>-C<sub>4</sub>)alkyl in the definition of R<sup>6</sup> and said (C<sub>1</sub>-C<sub>4</sub>)alkoxy in the definition of R<sup>6</sup> are optionally and independently substituted with up to five fluoro; and R<sup>19</sup> and R<sup>20</sup> are each independently (C<sub>1</sub>-C<sub>4</sub>)alkyl; or G<sup>3</sup>, G<sup>4</sup> and G<sup>5</sup> are taken separately and are each hydrogen; r is 1; R<sup>18</sup> is hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl

or phenyl optionally independently substituted with up to three hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl or (C<sub>1</sub>-C<sub>4</sub>)alkoxy, wherein said (C<sub>1</sub>-C<sub>4</sub>)alkyl in the definition of R<sup>6</sup> and said (C<sub>1</sub>-C<sub>4</sub>)alkoxy in the definition of R<sup>6</sup> are optionally and independently substituted with up to five fluoro; and

5 R<sup>19</sup> and R<sup>20</sup> are each independently hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl; or G<sup>3</sup> and G<sup>4</sup> are taken together and are (C<sub>1</sub>-C<sub>3</sub>)alkylene; r is 0 or 1; and R<sup>18</sup>, R<sup>19</sup>, R<sup>20</sup> and G<sup>5</sup> are hydrogen; or G<sup>4</sup> and G<sup>5</sup> are taken together and are (C<sub>1</sub>-C<sub>3</sub>)alkylene; r is 0 or 1; and R<sup>18</sup>, R<sup>19</sup>, R<sup>20</sup> and G<sup>3</sup> are hydrogen;

10 R<sup>17</sup> is SO<sub>2</sub>NR<sup>21</sup>R<sup>22</sup>, CONR<sup>21</sup>R<sup>22</sup>, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl, Ar<sup>2</sup>-carbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfinyl, Ar<sup>2</sup>-sulfonyl, Ar<sup>2</sup>-sufinyl and (C<sub>1</sub>-C<sub>6</sub>)alkyl; R<sup>21</sup> and R<sup>22</sup> are taken separately and are each independently selected from hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl and Ar<sup>2</sup>-(C<sub>0</sub>-C<sub>4</sub>)alkylenyl; or

15 R<sup>21</sup> and R<sup>22</sup> are taken together with the nitrogen atom to which they are attached to form azetidiny, pyrrolidiny, piperidiny, piperaziny, morpholinyl, azepiny, azabicyclo[3.2.2]nonany, azabicyclo[2.2.1]heptyl, 6,7-dihydro-5H-dibenzo[c,e]azepiny, 1,2,3,4-tetrahydro-isoquinoly or 5,6,7,8-tetrahydropyrido[4,3-d]pyrimidyl; said azetidiny in the definition of R<sup>21</sup> and R<sup>22</sup> is optionally substituted

20 independently with one substituent selected from hydroxy, amino, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said pyrrolidiny, piperidiny, azepiny in the definition of R<sup>21</sup> and R<sup>22</sup> are optionally substituted independently with up to two substituents independently selected from hydroxy,

25 amino, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said morpholinyl in the definition of R<sup>21</sup> and R<sup>22</sup> is optionally substituted with up to two substituents independently selected from hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and

30 (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said piperaziny in the definition of R<sup>21</sup> and R<sup>22</sup> is optionally substituted independently with up to three substituents independently selected from phenyl, pyridyl, pyrimidyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl and (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro; said

35 1,2,3,4-tetrahydro-isoquinoly and said 5,6,7,8-tetrahydropyrido[4,3-d]pyrimidyl in the definition of R<sup>21</sup> and R<sup>22</sup> are optionally substituted independently with up to three

substituents independently selected from hydroxy, amino, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; and said 6,7-dihydro-5H-dibenzo[c,e]azepinyl in the definition of R<sup>21</sup> and R<sup>22</sup> is optionally substituted with up to four substituents independently selected from hydroxy, amino, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said pyrimidyl, pyridyl and phenyl which are optionally substituted on said piperazine in the definition of R<sup>21</sup> and R<sup>22</sup> is optionally substituted with up to three substituents selected from hydroxy, amino, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro;

Ar<sup>2</sup> is independently defined as set forth for Ar and Ar<sup>1</sup> above;

said Ar<sup>2</sup> is optionally independently substituted as set forth for Ar and Ar<sup>1</sup> above;

R<sup>23</sup> is CONR<sup>25</sup>R<sup>26</sup> or SO<sub>2</sub>R<sup>25</sup>R<sup>26</sup>, wherein R<sup>25</sup> is hydrogen (C<sub>1</sub>-C<sub>4</sub>)alkyl or Ar<sup>3</sup>-(C<sub>0</sub>-C<sub>4</sub>)alkylenyl and R<sup>26</sup> is Ar<sup>3</sup>-(C<sub>0</sub>-C<sub>4</sub>)alkylenyl; provided that when Ar<sup>3</sup> is phenyl, naphthyl or biphenyl, then R<sup>23</sup> cannot be CONR<sup>25</sup>R<sup>26</sup> where R<sup>25</sup> is hydrogen or Ar<sup>3</sup> and R<sup>26</sup> is Ar<sup>3</sup>;

R<sup>24</sup> is hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl or phenyl optionally independently substituted with up to three hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl or (C<sub>1</sub>-C<sub>4</sub>)alkoxy, wherein said (C<sub>1</sub>-C<sub>4</sub>)alkyl in the definition of R<sup>6</sup> and said (C<sub>1</sub>-C<sub>4</sub>)alkoxy in the definition of R<sup>6</sup> are optionally and independently substituted with up to five fluoro; Ar<sup>3</sup> is independently defined as set forth for Ar and Ar<sup>1</sup> above;

said Ar<sup>3</sup> is optionally independently substituted as set forth for Ar and Ar<sup>1</sup> above;

R<sup>27</sup> is hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl;

R<sup>28</sup> and R<sup>29</sup> are each independently hydrogen, hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro, (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro, phenyl, pyridyl, pyrimidyl,

thienyl, furanyl, thiazolyl, oxazolyl, phenoxy, thiophenoxy, SO<sub>2</sub>NR<sup>30</sup>R<sup>31</sup>, CONR<sup>30</sup>R<sup>31</sup> or NR<sup>30</sup>R<sup>31</sup>; said thienyl, pyrimidyl, furanyl, thiazolyl and oxazolyl in the definition of R<sup>28</sup> and R<sup>29</sup> are optionally substituted by up to two hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said phenyl, pyridyl,

phenoxy and thiophenoxy in the definition of R<sup>28</sup> and R<sup>29</sup> are optionally substituted by

up to three hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro;

R<sup>30</sup> and R<sup>31</sup> are each independently hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or

5 phenyl, said phenyl is optionally substituted with up to three hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; or

R<sup>30</sup> and R<sup>31</sup> are taken together with the nitrogen to which they are attached to form indoliny, pyrrolidiny, piperidiny, piperaziny or morpholiny; said pyrrolidiny and

10 piperidiny in the definition of R<sup>30</sup> and R<sup>31</sup> are optionally substituted with up to two hydroxy, amino, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said indoliny and piperaziny in the definition of R<sup>30</sup> and R<sup>31</sup> are

15 optionally substituted with up to three hydroxy, amino, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said morpholiny in the definition of R<sup>30</sup> and R<sup>31</sup> is optionally substituted with up to two substituents independently selected from hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy

20 optionally substituted with up to five fluoro;

A is N optionally substituted with hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl and B is carbonyl; or

A is carbonyl and B is N optionally substituted with hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl;

R<sup>32</sup> is hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl;

R<sup>33</sup> is phenyl, pyridyl, pyrimidyl, thiazolyl, oxazolyl, benzyl, quinolyl, isoquinolyl,

25 phthaliziny, quinoxanly, benzothiazoyl, benzoxazolyl, benzofuranyl or benzothiényl;

said phenyl, pyridyl, pyrimidyl, thiazolyl, oxazolyl, benzyl, quinolyl, isoquinolyl,

phthaliziny, quinoxanly, benzothiazoyl, benzoxazolyl, benzofuranyl and benzothiényl in the definition of R<sup>33</sup> are optionally substituted with up to three phenyl, phenoxy,

NR<sup>34</sup>R<sup>35</sup>, halo, hydroxy, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl

30 optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro;

R<sup>34</sup> and R<sup>35</sup> are each independently hydrogen, (C<sub>1</sub>-C<sub>4</sub> alkyl), phenyl or

phenylsulfonyl; said phenyl and phenylsulfonyl in the definition of R<sup>34</sup> and R<sup>35</sup> are optionally substituted with up to three halo, hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally

substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro;

D is CO, CHOH or CH<sub>2</sub>;

E is O, NH or S;

5 R<sup>36</sup> and R<sup>37</sup> are taken separately and are each independently hydrogen, halo, cyano, hydroxy, amino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino, di-(C<sub>1</sub>-C<sub>6</sub>)alkylamino, pyrrolidino, piperidino, morpholino, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, Ar<sup>4</sup>, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro;

10 R<sup>38</sup>, R<sup>39</sup> and R<sup>40</sup> are each independently hydrogen or (C<sub>1</sub>-C<sub>4</sub>)-alkyl; Ar<sup>4</sup> is phenyl, furanyl, thienyl, pyridyl, pyrimidyl, pyrazinyl or pyridazinyl; said Ar<sup>4</sup> being optionally substituted with up to three hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; or

15 R<sup>36</sup> and R<sup>37</sup> are taken together on adjacent carbon atoms and are -O-(CH<sub>2</sub>)<sub>t</sub>-O-; t is 1, 2 or 3;

Y is (C<sub>2</sub>-C<sub>6</sub>)alkylene;

R<sup>44</sup>, R<sup>45</sup> and R<sup>46</sup> are each independently hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl;

m and n are each independently 1, 2 or 3, provided that the sum of m and n is 2, 3 or

20 4;

k is 0, 1, 2, 3 or 4;

Y<sup>1</sup> is a covalent bond, carbonyl, sulfonyl or oxycarbonyl;

R<sup>43</sup> is (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, Ar<sup>5</sup>-(C<sub>0</sub>-C<sub>4</sub>)alkylenyl, NR<sup>47</sup>R<sup>48</sup> or (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted with one to five fluoro; provided that when Y<sup>1</sup> is a covalent bond or

25 oxycarbonyl, then R<sup>43</sup> is not NR<sup>47</sup>R<sup>48</sup>;

R<sup>47</sup> and R<sup>48</sup> are taken separately and are each independently selected from hydrogen, Ar<sup>5</sup>, (C<sub>1</sub>-C<sub>6</sub>)alkyl and Ar<sup>5</sup>-(C<sub>0</sub>-C<sub>4</sub>)alkylenyl; or

R<sup>47</sup> and R<sup>48</sup> are taken together with the nitrogen atom to which they are attached to form azetidiny, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, azepinyl,

30 azabicyclo[3.2.2]nonanyl, azabicyclo[2.2.1]heptyl, 1,2,3,4-tetrahydroisoquinolyl, 6,7-dihydro-5H-dibenzo[c,e]azepinyl or 5,6,7,8-tetrahydropyrido[4,3-d]pyrimidyl; said azetidiny in the definition of R<sup>47</sup> and R<sup>48</sup> are optionally substituted with one hydroxy,

amino, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five

35 fluoro; said pyrrolidinyl, piperidinyl and azepinyl in the definition of R<sup>47</sup> and R<sup>48</sup> are



optionally substituted with up to two hydroxy, amino, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said morpholinyl in the definition of R<sup>47</sup> and R<sup>48</sup> is optionally substituted with up to two substituents

- 5 independently selected from hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said piperazinyl, 1,2,3,4-tetrahydroisoquinolyl and 5,6,7,8-tetrahydro[4,3-d]pyrimidyl in the definition of R<sup>47</sup> and R<sup>48</sup> are optionally substituted with up to three hydroxy, amino, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; and said 6,7-dihydro-5H-dibenzo[c,e]azepinyl in the definition of R<sup>47</sup> and R<sup>48</sup> are optionally substituted with up to four hydroxy, amino, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro;
- 10 substituted with up to five fluoro;
- 15 substituted with up to five fluoro;

Ar<sup>5</sup> is independently defined as set forth for Ar and Ar<sup>1</sup> above;

Ar<sup>5</sup> is optionally independently substituted as set forth for Ar and Ar<sup>1</sup> above;

R<sup>42</sup> and R<sup>42a</sup> are independently hydrogen, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, Ar<sup>6</sup>-(C<sub>0</sub>-C<sub>3</sub>)alkylenyl, Ar<sup>6</sup>-(C<sub>2</sub>-C<sub>4</sub>)alkenyl, Ar<sup>6</sup>-carbonyl or (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted with up to five fluoro;

20 fluoro;

Ar<sup>6</sup> is independently defined as set forth for Ar and Ar<sup>1</sup> above;

Ar<sup>6</sup> is optionally independently substituted as set forth for Ar and Ar<sup>1</sup> above; and

R<sup>41</sup> and R<sup>41a</sup> are each independently hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl.

166. A compound of claim 165 selected from 1R-(4-{4-[2-(1R-butyryloxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl}-pyrimidin-2-yl)-ethyl butyrate; 1R-(4-{4-[2-(1S-butyryloxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl}-pyrimidin-2-yl)-ethyl butyrate; 1S-(4-{4-[2-(1R-butyryloxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl}-pyrimidin-2-yl)-ethyl butyrate; (E)-1R-{4-[4-(2-methyl-32-phenyl-acryloyl)-piperazin-1-yl]-pyrimidin-2-yl}-ethyl acetate; (R)-1-[4-(4-quinoxalin-2-yl-piperazin-1-yl)-pyrimidin-2-yl]-ethyl acetate; 1R-(4-{4-[2-(1RS-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl}-pyrimidin-2-yl)-ethyl butyrate; 1RS-(4-{4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-3R,5S-dimethyl-piperazin-1-yl}-pyrimidin-2-yl)-ethyl butyrate; 1R-[4-(3S-methyl-4-oxazolo[5,4-b]pyridin-2-yl-piperazin-1-yl)-pyrimidin-2-yl]-ethyl butyrate; 1R-(4-[3R,5S-dimethyl-4-(4-methyl-6-phenyl-[1,3,5]triazin-2-yl)-piperazin-1-yl]-pyrimidin-2-yl)-ethyl butyrate; 1R-(4-[4-(4-


25 ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl)-pyrimidin-2-yl)-ethyl butyrate;

30 ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl)-pyrimidin-2-yl)-ethyl butyrate; 1S-(4-{4-[2-(1R-butyryloxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl}-pyrimidin-2-yl)-ethyl butyrate; (E)-1R-{4-[4-(2-methyl-32-phenyl-acryloyl)-piperazin-1-yl]-pyrimidin-2-yl}-ethyl acetate; (R)-1-[4-(4-quinoxalin-2-yl-piperazin-1-yl)-pyrimidin-2-yl]-ethyl acetate; 1R-(4-{4-[2-(1RS-hydroxy-ethyl)-pyrimidin-4-yl]-2R,6S-dimethyl-piperazin-1-yl}-pyrimidin-2-yl)-ethyl butyrate; 1RS-(4-{4-[2-(1R-hydroxy-ethyl)-pyrimidin-4-yl]-3R,5S-dimethyl-piperazin-1-yl}-pyrimidin-2-yl)-ethyl butyrate; 1R-[4-(3S-methyl-4-oxazolo[5,4-b]pyridin-2-yl-piperazin-1-yl)-pyrimidin-2-yl]-ethyl butyrate; 1R-(4-[3R,5S-dimethyl-4-(4-methyl-6-phenyl-[1,3,5]triazin-2-yl)-piperazin-1-yl]-pyrimidin-2-yl)-ethyl butyrate; 1R-(4-[4-(4-

35 [1,3,5]triazin-2-yl)-piperazin-1-yl]-pyrimidin-2-yl)-ethyl butyrate; 1R-(4-[4-(4-

cyclopropyl-[1,3,5]triazin-2-yl)-3R,5S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethyl butyrate; 1R-{4-[4-(4-cyclopropyl-[1,3,5]triazin-2-yl)-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethyl butyrate; 1R-{4-[4-(4,6-dimethyl-[1,3,5]triazin-2-yl)-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethyl butyrate; 1R-{4-[4-(4-hydroxymethyl-6-phenyl-[1,3,5]triazin-2-yl)-2R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethyl butyrate; 1R-{4-[4-(4-methoxy-6-methoxymethyl-[1,3,5]triazin-2-yl)-R,6S-dimethyl-piperazin-1-yl]-pyrimidin-2-yl}-ethyl butyrate; and 1R-{4-[2R,6S-dimethyl-4-(4-methyl-[1,3,5]triazin-2-yl)-piperazin-1-yl]-pyrimidin-2-yl}-ethyl butyrate.

167. A compound of the formula I<sup>B</sup>,









wherein:


R<sup>1</sup> is C-(OR<sup>81</sup>)R<sup>4</sup>R<sup>5</sup>, where R<sup>81</sup> is an acyl radical of a carboxylic acid aldose reductase inhibitor;

R<sup>4</sup> and R<sup>5</sup> are each independently hydrogen, methyl, ethyl or hydroxy-(C<sub>1</sub>-C<sub>3</sub>)alkyl;

R<sup>2</sup> is hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl or (C<sub>1</sub>-C<sub>4</sub>)alkoxy;

R<sup>3</sup> is a radical of the formula





said radical of formula  $R^{3b}$  is substituted by  $R^{18}$ ,  $R^{19}$  and  $R^{20}$ ;

G,  $G^1$  and  $G^2$  are taken separately and are each hydrogen and  $R^6$  is hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl or phenyl optionally independently substituted with up to three hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl or (C<sub>1</sub>-C<sub>4</sub>)alkoxy, wherein said (C<sub>1</sub>-C<sub>4</sub>)alkyl in the definition of  $R^6$  and said (C<sub>1</sub>-C<sub>4</sub>)alkoxy in the definition of  $R^6$  are optionally and independently substituted with up to five fluoro;  $R^7$  and  $R^8$  are each independently hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl; or

G and  $G^1$  are taken together and are (C<sub>1</sub>-C<sub>3</sub>)alkylene and  $R^6$ ,  $R^7$ ,  $R^8$  and  $G^2$  are hydrogen; or

$G^1$  and  $G^2$  are taken together and are (C<sub>1</sub>-C<sub>3</sub>)alkylene and  $R^6$ ,  $R^7$ ,  $R^8$  and G are hydrogen;

q is 0 or 1;

X is a covalent bond, -(C=NR<sup>10</sup>)-, oxycarbonyl, vinylenylcarbonyl, oxy(C<sub>1</sub>-

C<sub>4</sub>)alkylenylcarbonyl, (C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl, (C<sub>3</sub>-C<sub>4</sub>)alkenylcarbonyl, thio(C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl, vinylenylsulfonyl, sulfinyl-(C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl, sulfonyl-(C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl or carbonyl(C<sub>0</sub>-C<sub>4</sub>)alkylenylcarbonyl; wherein said oxy(C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl, (C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl, (C<sub>3</sub>-C<sub>4</sub>)alkenylcarbonyl and thio(C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl in the definition of X are each optionally and independently substituted with up to two (C<sub>1</sub>-C<sub>4</sub>)alkyl, benzyl or Ar; said vinylenylsulfonyl and said vinylenylcarbonyl in the definition of X are optionally substituted independently on one or two vinylenyl carbons with (C<sub>1</sub>-C<sub>4</sub>)alkyl, benzyl or Ar; and said carbonyl(C<sub>0</sub>-C<sub>4</sub>)alkylenylcarbonyl in the definition of X is optionally substituted independently with up to three (C<sub>1</sub>-C<sub>4</sub>)alkyl, benzyl or Ar;

$R^{10}$  is hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl;

$R^9$  is (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, Ar<sup>1</sup>-(C<sub>0</sub>-C<sub>3</sub>)alkylenyl or (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted with up to five fluoro; provided that when q = 0 and X is a covalent bond, oxycarbonyl or (C<sub>1</sub>-C<sub>4</sub>)alkylenylcarbonyl, then  $R^9$  is not (C<sub>1</sub>-C<sub>6</sub>)alkyl;

Ar and Ar<sup>1</sup> are independently a fully saturated, partially saturated or fully unsaturated five- to eight-membered ring optionally having up to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of two fused independently partially saturated, fully saturated or fully unsaturated five- to seven-membered rings, taken independently, optionally having up to four heteroatoms selected independently from nitrogen, sulfur and oxygen, or a tricyclic ring consisting of three fused independently partially saturated, fully saturated or fully

unsaturated five to seven membered rings, taken independently, optionally having up to four heteroatoms selected independently from nitrogen, sulfur and oxygen, said partially saturated, fully saturated ring or fully unsaturated monocyclic ring, bicyclic ring or tricyclic ring optionally having one or two oxo groups substituted on carbon or one or two oxo groups substituted on sulfur;

Ar and Ar<sup>1</sup> are optionally independently substituted on carbon or nitrogen, on one ring if the moiety is monocyclic, on one or both rings if the moiety is bicyclic, or on one, two or three rings if the moiety is tricyclic, with up to a total of four substituents independently selected from R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup>; wherein R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> are

each taken separately and are each independently halo, formyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylenyloxycarbonyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, C(OH)R<sup>15</sup>R<sup>16</sup>, naphthyl, phenyl, imidazolyl, pyridyl, triazolyl, morpholinyl, (C<sub>0</sub>-C<sub>4</sub>)alkylsulfamoyl, N-(C<sub>0</sub>-C<sub>4</sub>)alkylcarbamoyl, N,N-di-(C<sub>1</sub>-C<sub>4</sub>)alkylcarbamoyl, N-phenylcarbamoyl, N-(C<sub>1</sub>-C<sub>4</sub>)alkyl-N-phenylcarbamoyl, N,N-diphenyl carbamoyl, (C<sub>1</sub>-C<sub>4</sub>)alkylcarbonylamido, (C<sub>3</sub>-C<sub>7</sub>)cycloalkylcarbonylamido, phenylcarbonylamido, piperidinyl, pyrrolidinyl, piperazinyl, cyano, benzimidazolyl, amino, anilino, pyrimidyl, oxazolyl, isoxazolyl, tetrazolyl, thienyl, thiazolyl, benzothiazolyl, pyrrolyl, pyrazolyl, tetrahydroquinolyl, tetrahydroisoquinolyl, benzoxazolyl, pyridazinyl, pyridyloxy, pyridylsulfanyl, furanyl, 8-(C<sub>1</sub>-C<sub>4</sub>)alkyl-3,8-diaza[3.2.1]bicyclooctyl, 3,5-dioxo-1,2,4-triazinyl, phenoxy, thiophenoxy, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfanyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said naphthyl, phenyl, pyridyl, piperidinyl, benzimidazolyl, pyrimidyl, thienyl, benzothiazolyl, pyrrolyl, tetrahydroquinolyl, tetrahydroisoquinolyl, benzoxazolyl, pyridazinyl, pyridyloxy, pyridylsulfanyl, furanyl, thiophenoxy, anilino and phenoxy in the definition of R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> are optionally substituted with up to three substituents independently selected from hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said imidazolyl, oxazolyl, isoxazolyl, thiazolyl and pyrazolyl in the definition of R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> are optionally substituted with up to two substituents independently selected from hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said morpholinyl in the definition of R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> is optionally substituted with up to two substituents independently selected from (C<sub>1</sub>-C<sub>4</sub>)alkyl; said pyrrolidinyl in the definition of R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> is

optionally substituted with up to two substituents independently selected from hydroxy, hydroxy-(C<sub>1</sub>-C<sub>3</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said piperazinyl in the definition of R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> is optionally substituted with up to three substituents independently selected from (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, hydroxy-(C<sub>1</sub>-C<sub>3</sub>)alkyl, phenyl, pyridyl, (C<sub>0</sub>-C<sub>4</sub>)alkylsulfamoyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said triazolyl in the definition of R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> is optionally substituted with hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said tetrazolyl in the definition of R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> is optionally substituted with hydroxy-(C<sub>2</sub>-C<sub>3</sub>)alkyl or (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro; and said phenyl and pyridyl which are optionally substituted on piperazine in the definition of R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> are optionally substituted with up to three hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; or R<sup>11</sup> and R<sup>12</sup> are taken together on adjacent carbon atoms and are -CH<sub>2</sub>OC(CH<sub>3</sub>)<sub>2</sub>OCH<sub>2</sub>- or -O-(CH<sub>2</sub>)<sub>p</sub>-O-, and R<sup>13</sup> and R<sup>14</sup> are taken separately and are each independently hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl; p is 1, 2 or 3; R<sup>15</sup> and R<sup>16</sup> are taken separately and are each independently hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro; or R<sup>15</sup> and R<sup>16</sup> are taken separately and R<sup>15</sup> is hydrogen and R<sup>16</sup> is (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, hydroxy-(C<sub>1</sub>-C<sub>3</sub>)alkyl, phenyl, pyridyl, pyrimidyl, thienyl, furanyl, thiazolyl, oxazolyl, imidazolyl, benzothiazolyl or benzoxazolyl; or R<sup>15</sup> and R<sup>16</sup> are taken together and are (C<sub>3</sub>-C<sub>6</sub>)alkylene; G<sup>3</sup>, G<sup>4</sup> and G<sup>5</sup> are taken separately and are each hydrogen; r is 0; R<sup>18</sup> is hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl or phenyl optionally independently substituted with up to three hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl or (C<sub>1</sub>-C<sub>4</sub>)alkoxy, wherein said (C<sub>1</sub>-C<sub>4</sub>)alkyl in the definition of R<sup>6</sup> and said (C<sub>1</sub>-C<sub>4</sub>)alkoxy in the definition of R<sup>6</sup> are optionally and independently substituted with up to five fluoro; and R<sup>19</sup> and R<sup>20</sup> are each independently (C<sub>1</sub>-C<sub>4</sub>)alkyl; or G<sup>3</sup>, G<sup>4</sup> and G<sup>5</sup> are taken separately and are each hydrogen; r is 1; R<sup>18</sup> is hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl

or phenyl optionally independently substituted with up to three hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl or (C<sub>1</sub>-C<sub>4</sub>)alkoxy, wherein said (C<sub>1</sub>-C<sub>4</sub>)alkyl in the definition of R<sup>6</sup> and said (C<sub>1</sub>-C<sub>4</sub>)alkoxy in the definition of R<sup>6</sup> are optionally and independently substituted with up to five fluoro; and

5 R<sup>19</sup> and R<sup>20</sup> are each independently hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl; or

G<sup>3</sup> and G<sup>4</sup> are taken together and are (C<sub>1</sub>-C<sub>3</sub>)alkylene; r is 0 or 1; and R<sup>18</sup>, R<sup>19</sup>, R<sup>20</sup> and G<sup>5</sup> are hydrogen; or

G<sup>4</sup> and G<sup>5</sup> are taken together and are (C<sub>1</sub>-C<sub>3</sub>)alkylene; r is 0 or 1; and R<sup>18</sup>, R<sup>19</sup>, R<sup>20</sup> and G<sup>3</sup> are hydrogen;

10 R<sup>17</sup> is SO<sub>2</sub>NR<sup>21</sup>R<sup>22</sup>, CONR<sup>21</sup>R<sup>22</sup>, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylcarbonyl, Ar<sup>2</sup>-carbonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfinyl, Ar<sup>2</sup>-sulfonyl, Ar<sup>2</sup>-sufinyl and (C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sup>21</sup> and R<sup>22</sup> are taken separately and are each independently selected from hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl and Ar<sup>2</sup>-(C<sub>6</sub>-C<sub>4</sub>)alkylenyl; or

15 R<sup>21</sup> and R<sup>22</sup> are taken together with the nitrogen atom to which they are attached to form azetidiny, pyrrolidiny, piperidiny, piperaziny, morpholiny, azepiny, azabicyclo[3.2.2]nonany, azabicyclo[2.2.1]heptyl, 6,7-dihydro-5H-dibenzo[c,e]azepiny, 1,2,3,4-tetrahydro-isoquinoly or 5,6,7,8-tetrahydropyrido[4,3-d]pyrimidy; said azetidiny in the definition of R<sup>21</sup> and R<sup>22</sup> is optionally substituted

20 independently with one substituent selected from hydroxy, amino, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said pyrrolidiny, piperidiny, azepiny in the definition of R<sup>21</sup> and R<sup>22</sup> are optionally substituted independently with up to two substituents independently selected from hydroxy,

25 amino, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said morpholiny in the definition of R<sup>21</sup> and R<sup>22</sup> is optionally substituted with up to two substituents independently selected from hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and

30 (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said piperaziny in the definition of R<sup>21</sup> and R<sup>22</sup> is optionally substituted independently with up to three substituents independently selected from phenyl, pyridyl, pyrimidy, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl and (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro; said 1,2,3,4-tetrahydro-isoquinoly and said 5,6,7,8-tetrahydropyrido[4,3-d]pyrimidy in the

35 definition of R<sup>21</sup> and R<sup>22</sup> are optionally substituted independently with up to three



substituents independently selected from hydroxy, amino, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; and said 6,7-dihydro-5H-dibenzo[c,e]azepinyl in the definition of R<sup>21</sup> and R<sup>22</sup> is optionally substituted with up to four substituents independently selected from hydroxy, amino, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said pyrimidyl, pyridyl and phenyl which are optionally substituted on said piperazine in the definition of R<sup>21</sup> and R<sup>22</sup> is optionally substituted with up to three substituents selected from hydroxy, amino, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro;

Ar<sup>2</sup> is independently defined as set forth for Ar and Ar<sup>1</sup> above;

said Ar<sup>2</sup> is optionally independently substituted as set forth for Ar and Ar<sup>1</sup> above;

R<sup>23</sup> is CONR<sup>25</sup>R<sup>26</sup> or SO<sub>2</sub>R<sup>25</sup>R<sup>26</sup>, wherein R<sup>25</sup> is hydrogen (C<sub>1</sub>-C<sub>4</sub>)alkyl or Ar<sup>3</sup>-(C<sub>0</sub>-C<sub>4</sub>)alkylenyl and R<sup>26</sup> is Ar<sup>3</sup>-(C<sub>0</sub>-C<sub>4</sub>)alkylenyl; provided that when Ar<sup>3</sup> is phenyl, naphthyl or biphenyl, then R<sup>23</sup> cannot be CONR<sup>25</sup>R<sup>26</sup> where R<sup>25</sup> is hydrogen or Ar<sup>3</sup> and R<sup>26</sup> is Ar<sup>3</sup>;

R<sup>24</sup> is hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl or phenyl optionally independently substituted with up to three hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl or (C<sub>1</sub>-C<sub>4</sub>)alkoxy, wherein said (C<sub>1</sub>-C<sub>4</sub>)alkyl in the definition of R<sup>6</sup> and said (C<sub>1</sub>-C<sub>4</sub>)alkoxy in the definition of R<sup>6</sup> are optionally and independently substituted with up to five fluoro; Ar<sup>3</sup> is independently defined as set forth for Ar and Ar<sup>1</sup> above;

said Ar<sup>3</sup> is optionally independently substituted as set forth for Ar and Ar<sup>1</sup> above;

R<sup>27</sup> is hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl;

R<sup>28</sup> and R<sup>29</sup> are each independently hydrogen, hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro, (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro, phenyl, pyridyl, pyrimidyl, thienyl, furanyl, thiazolyl, oxazolyl, phenoxy, thiophenoxy, SO<sub>2</sub>NR<sup>30</sup>R<sup>31</sup>, CONR<sup>30</sup>R<sup>31</sup> or NR<sup>30</sup>R<sup>31</sup>; said thienyl, pyrimidyl, furanyl, thiazolyl and oxazolyl in the definition of R<sup>28</sup> and R<sup>29</sup> are optionally substituted by up to two hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said phenyl, pyridyl, phenoxy and thiophenoxy in the definition of R<sup>28</sup> and R<sup>29</sup> are optionally substituted by

up to three hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro;

R<sup>30</sup> and R<sup>31</sup> are each independently hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or

5 phenyl, said phenyl is optionally substituted with up to three hydroxy, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; or

R<sup>30</sup> and R<sup>31</sup> are taken together with the nitrogen to which they are attached to form indolinyl, pyrrolidinyl, piperidinyl, piperazinyl or morpholinyl; said pyrrolidinyl and

10 piperidinyl in the definition of R<sup>30</sup> and R<sup>31</sup> are optionally substituted with up to two hydroxy, amino, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said indolinyl and piperazinyl in the definition of R<sup>30</sup> and R<sup>31</sup> are

optionally substituted with up to three hydroxy, amino, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said morpholinyl in the definition of R<sup>30</sup> and R<sup>31</sup> is optionally substituted with up to two substituents independently selected from hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro;

A is N optionally substituted with hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl and B is carbonyl; or

A is carbonyl and B is N optionally substituted with hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl;

R<sup>32</sup> is hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl;

R<sup>33</sup> is phenyl, pyridyl, pyrimidyl, thiazolyl, oxazolyl, benzyl, quinolyl, isoquinolyl,

25 phthalizinyl, quinoxanlyl, benzothiazoyl, benzoxazolyl, benzofuranyl or benzothienyl; said phenyl, pyridyl, pyrimidyl, thiazolyl, oxazolyl, benzyl, quinolyl, isoquinolyl, phthalizinyl, quinoxanlyl, benzothiazoyl, benzoxazolyl, benzofuranyl and benzothienyl in the definition of R<sup>33</sup> are optionally substituted with up to three phenyl, phenoxy,

NR<sup>34</sup>R<sup>35</sup>, halo, hydroxy, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro;

R<sup>34</sup> and R<sup>35</sup> are each independently hydrogen, (C<sub>1</sub>-C<sub>4</sub> alkyl), phenyl or

phenylsulfonyl; said phenyl and phenylsulfonyl in the definition of R<sup>34</sup> and R<sup>35</sup> are optionally substituted with up to three halo, hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally

substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro;

D is CO, CHOH or CH<sub>2</sub>;

E is O, NH or S;

5 R<sup>36</sup> and R<sup>37</sup> are taken separately and are each independently hydrogen, halo, cyano, hydroxy, amino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino, di-(C<sub>1</sub>-C<sub>6</sub>)alkylamino, pyrrolidino, piperidino, morpholino, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, Ar<sup>4</sup>, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro;

10 R<sup>38</sup>, R<sup>39</sup> and R<sup>40</sup> are each independently hydrogen or (C<sub>1</sub>-C<sub>4</sub>)-alkyl; Ar<sup>4</sup> is phenyl, furanyl, thienyl, pyridyl, pyrimidyl, pyrazinyl or pyridazinyl; said Ar<sup>4</sup> being optionally substituted with up to three hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; or

15 R<sup>36</sup> and R<sup>37</sup> are taken together on adjacent carbon atoms and are -O-(CH<sub>2</sub>)<sub>t</sub>-O-; t is 1, 2 or 3;

Y is (C<sub>2</sub>-C<sub>6</sub>)alkylene;

R<sup>44</sup>, R<sup>45</sup> and R<sup>46</sup> are each independently hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl;

m and n are each independently 1, 2 or 3, provided that the sum of m and n is 2, 3 or 4;

20 k is 0, 1, 2, 3 or 4;

Y<sup>1</sup> is a covalent bond, carbonyl, sulfonyl or oxycarbonyl;

R<sup>43</sup> is (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, Ar<sup>5</sup>-(C<sub>0</sub>-C<sub>4</sub>)alkylenyl, NR<sup>47</sup>R<sup>48</sup> or (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted with one to five fluoro; provided that when Y<sup>1</sup> is a covalent bond or 25 oxycarbonyl, then R<sup>43</sup> is not NR<sup>47</sup>R<sup>48</sup>;

R<sup>47</sup> and R<sup>48</sup> are taken separately and are each independently selected from hydrogen, Ar<sup>5</sup>, (C<sub>1</sub>-C<sub>6</sub>)alkyl and Ar<sup>5</sup>-(C<sub>0</sub>-C<sub>4</sub>)alkylenyl; or

R<sup>47</sup> and R<sup>48</sup> are taken together with the nitrogen atom to which they are attached to form azetidiny, pyrrolidiny, piperidiny, piperazinyl, morpholinyl, azepiny, 30 azabicyclo[3.2.2]nonanyl, azabicyclo[2.2.1]heptyl, 1,2,3,4-tetrahydroisoquinolyl, 6,7-dihydro-5H-dibenzo[c,e]azepiny or 5,6,7,8-tetrahydropyrido[4,3-d]pyrimidyl; said azetidiny in the definition of R<sup>47</sup> and R<sup>48</sup> are optionally substituted with one hydroxy, amino, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five 35 fluoro; said pyrrolidiny, piperidiny and azepiny in the definition of R<sup>47</sup> and R<sup>48</sup> are

optionally substituted with up to two hydroxy, amino, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said morpholinyl in the definition of R<sup>47</sup> and R<sup>48</sup> is optionally substituted with up to two substituents

5 independently selected from hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro and (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; said piperazinyl, 1,2,3,4-tetrahydroisoquinolyl and 5,6,7,8-tetrahydro[4,3-d]pyrimidyl in the definition of R<sup>47</sup> and R<sup>48</sup> are optionally substituted with up to three hydroxy, amino, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro; and said 6,7-dihydro-5H-dibenzo[c,e]azepinyl in the definition of R<sup>47</sup> and R<sup>48</sup> are optionally substituted with up to four hydroxy, amino, halo, hydroxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with up to five fluoro or (C<sub>1</sub>-C<sub>4</sub>)alkoxy optionally substituted with up to five fluoro;

Ar<sup>5</sup> is independently defined as set forth for Ar and Ar<sup>1</sup> above;

Ar<sup>5</sup> is optionally independently substituted as set forth for Ar and Ar<sup>1</sup> above;

R<sup>42</sup> and R<sup>42a</sup> are independently hydrogen, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, Ar<sup>6</sup>-(C<sub>0</sub>-C<sub>3</sub>)alkylenyl, Ar<sup>6</sup>-(C<sub>2</sub>-C<sub>4</sub>)alkenyl, Ar<sup>6</sup>-carbonyl or (C<sub>1</sub>-C<sub>6</sub>)alkyl optionally substituted with up to five fluoro;

Ar<sup>6</sup> is independently defined as set forth for Ar and Ar<sup>1</sup> above;


Ar<sup>6</sup> is optionally independently substituted as set forth for Ar and Ar<sup>1</sup> above; and

R<sup>41</sup> and R<sup>41a</sup> are each independently hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl.

168. A compound of claim 167 wherein R<sup>81</sup> is the acyl radical of ponalrestat, tolrestat, zenarastat, zopolrestat, epalrestat, ZD5522 or sorbinil.


169. A compound selected from (E)-[4-oxo-3-(5-trifluoromethyl-benzothiazol-2-ylmethyl)-3,4-dihydro-phthalazin-1-yl]-acetic acid 1R-[4-(4-quinoxalin-2-yl-piperazin-1-yl)-pyrimidin-2-yl]-ethyl ester and (E)-[4-Oxo-3-(5-trifluoromethyl-benzothiazol-2-ylmethyl)-3,4-dihydro-phthalazin-1-yl]-acetic acid 1R-{4-[4-(3-thiophen-2-yl-acryloyl)-piperazin-1-yl]-pyrimidin-2-yl}-ethyl ester.

170. A compound of the formula Z



Z

171. A compound of the formula ZZ,




ZZ

wherein R<sup>100</sup> is (C<sub>1</sub>-C<sub>8</sub>)alkyl, benzyl or phenyl wherein said benzyl and phenyl are optionally substituted with up to three halo or (C<sub>1</sub>-C<sub>4</sub>)alkyl.

172. A compound of claim 171 wherein R<sup>100</sup> is (C<sub>1</sub>-C<sub>4</sub>)alkyl.

173. A compound of claim 172 wherein R<sup>100</sup> is n-butyl or ethyl.

174. A compound of the formula III,



III

wherein:

R<sup>100</sup> is (C<sub>1</sub>-C<sub>8</sub>)alkyl, benzyl or phenyl wherein said benzyl and phenyl are optionally substituted with up to three halo or (C<sub>1</sub>-C<sub>4</sub>)alkyl; and


R<sup>101</sup> is hydrogen or a suitable amine protecting group.

175. A compound of claim 174 wherein R<sup>100</sup> is (C<sub>1</sub>-C<sub>4</sub>)alkyl and R<sup>101</sup> is benzyl or tert-butyloxycarbonyl.

176. A compound of claim 175 wherein  $R^{100}$  is n-butyl or ethyl and  $R^{101}$  is benzyl.

177. A compound of claim 175 wherein  $R^{100}$  is n-butyl or ethyl and  $R^{101}$  is tert-butyloxycarbonyl.

5 178. A process for preparing a compound of the formula Z,



Z

comprising:

- a) reacting R-(+)-2-hydroxy-propionamide with triethyloxonium  
10 tetrafluoroborate in a reaction inert solvent for 10 minutes to 24 hours at 0 °C to ambient temperature to form the corresponding imidate;
- b) reacting said corresponding imidate with anhydrous ammonia in a reaction inert solvent for 2 hours to 24 hours at 0 °C to ambient temperature to form R-(+)-2-hydroxy-propionamidine hydrochloride; and
- 15 c) reacting said R-(+)-2-hydroxy-propionamidine hydrochloride with ethyl 3-hydroxy-acrylate sodium salt and a suitable base in a reaction inert solvent to form said compound of formula Z.

179. A pharmaceutical composition comprising a compound of claim 1, a prodrug thereof or a pharmaceutically acceptable salt of said prodrug or said  
20 compound, and a glycogen phosphorylase inhibitor (GPI), a prodrug of said GPI or a pharmaceutically acceptable salt of said GPI or said prodrug.

180. A kit comprising:

- a. a compound of claim 1, a prodrug thereof or a pharmaceutically acceptable salt of said prodrug or said compound in a first unit  
25 dosage form;
- b. a glycogen phosphorylase inhibitor (GPI), a prodrug thereof or a pharmaceutically acceptable salt of said prodrug or said GPI in a second unit dosage form; and
- c. a container.

181. A method of treating or preventing diabetic complications in a mammal comprising administering to said mammal a pharmaceutical composition of claim 179.

5 182. A method of treating hyperglycemia in a mammal comprising administering to said mammal a pharmaceutical composition of claim 179.

183. A method of treating ischemia in a mammal suffering from ischemia comprising administering to said mammal a pharmaceutical composition of claim 179.

10 184. A method of treating diabetes in a mammal comprising administering to said mammal a pharmaceutical composition of claim 179.

185. A method of treating diabetic complications in a mammal comprising administering to said mammal a compound of claim 1, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug and a glycogen phosphorylase inhibitor (GPI), a prodrug of said GPI or a pharmaceutically acceptable salt of said GPI or said prodrug.

15 186. A method of treating hyperglycemia in a mammal comprising administering to said mammal a compound of claim 1, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug and a glycogen phosphorylase inhibitor (GPI), a prodrug of said GPI or a pharmaceutically acceptable salt of said GPI or said prodrug.

20 187. A method of treating ischemia in a mammal comprising administering to said mammal a compound of claim 1, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug and a glycogen phosphorylase inhibitor (GPI), a prodrug of said GPI or a pharmaceutically acceptable salt of said GPI or said prodrug.

25 188. A method of treating diabetes in a mammal comprising administering to said mammal a compound of claim 1, a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug and a glycogen phosphorylase inhibitor (GPI), a prodrug of said GPI or a pharmaceutically acceptable salt of said GPI or said prodrug.

30